

Levetiracetam Extended-Release Tablets  
500 mg and 750 mg  
Rx Only  
Part 1 of 2

8183226 R7

## HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use LEVETIRACETAM EXTENDED-RELEASE TABLETS safely and effectively. See full prescribing information for LEVETIRACETAM EXTENDED-RELEASE TABLETS.

LEVETIRACETAM extended-release tablets, for oral use

Initial U.S. Approval: 1999

**INDICATIONS AND USAGE**  
Levetiracetam extended-release tablets are indicated for adjunctive therapy in the treatment of partial onset seizures in patients 12 years of age and older with epilepsy (1)

**DOSAGE AND ADMINISTRATION**  
Initiate treatment with a dose of 1000 mg once daily; increase by 1000 mg every 2 weeks to a maximum recommended dose of 3000 mg once daily (2)

See full prescribing information for use in patients with impaired renal function (2.1)

**DOSAGE FORMS AND STRENGTHS**

- 500 mg white, film-coated extended-release tablet (3)
- 750 mg white, film-coated extended-release tablet (3)

**CONTRAINDICATIONS**

None (4)

## FULL PRESCRIBING INFORMATION: CONTENTS\*

### 1 INDICATIONS AND USAGE

#### 2 DOSAGE AND ADMINISTRATION

- 2.1 Recommended Dosing
- 2.2 Dosage Adjustments in Adult Patients with Renal Impairment

#### 3 DOSAGE FORMS AND STRENGTHS

#### 4 CONTRAINDICATIONS

#### 5 WARNINGS AND PRECAUTIONS

- 5.1 Behavioral Abnormalities and Psychotic Symptoms
- 5.2 Suicidal Behavior and Ideation
- 5.3 Somnolence and Fatigue
- 5.4 Serious Dermatological Reactions
- 5.5 Coordination Difficulties
- 5.6 Withdrawal Seizures
- 5.7 Hematologic Abnormalities
- 5.8 Seizure Control During Pregnancy

#### 6 ADVERSE REACTIONS

- 6.1 Clinical Trials Experience
- 6.2 Postmarketing Experience

#### 8 USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy
- 8.2 Labor and Delivery
- 8.3 Nursing Mothers
- 8.4 Pediatric Use

Revised: 05/2016

## FULL PRESCRIBING INFORMATION

### 1 INDICATIONS AND USAGE

Levetiracetam extended-release tablets are indicated as adjunctive therapy in the treatment of partial onset seizures in patients 12 years of age and older with epilepsy.

#### 2 DOSAGE AND ADMINISTRATION

##### 2.1 Recommended Dosing

Levetiracetam extended-release tablets are administered once daily.

Initiate treatment with a dose of 1000 mg once daily. The once daily dosage may be adjusted in increments of 1000 mg every 2 weeks to a maximum recommended daily dose of 3000 mg/day.

##### 2.2 Dosage Adjustment in Adult Patients with Renal Impairment

Levetiracetam extended-release tablet dosing must be individualized according to the patient's renal function status. Recommended dosage adjustments for adults are shown in Table 1. In order to calculate the dose recommended for patients with renal impairment, creatinine clearance adjusted for body surface area must be calculated. To do this, an estimate of the patient's creatinine clearance (CL<sub>cr</sub>) in mL/min must first be calculated using the following formula:

$$CL_{cr} = \frac{[140 - \text{age (years)}] \times \text{weight (kg)}}{72 \times \text{serum creatinine (mg/dL)}} \times (0.85 \text{ for female patients})$$

Then CL<sub>cr</sub> is adjusted for body surface area (BSA) as follows:

$$CL_{cr} \text{ (mL/min)} = \frac{CL_{cr} \text{ (mL/min)}}{BSA \text{ subject (m}^2)} \times 1.73$$

Table 1. Dosage Adjustment Regimen for Adult Patients With Renal Impairment

| Group    | Creatinine Clearance (mL/min/1.73m <sup>2</sup> ) | Dosage (mg)  | Frequency      |
|----------|---|--------------|----------------|
| Normal   | >80   | 1000 to 3000 | Every 24 hours |
| Mild     | 50–80   | 1000 to 2000 | Every 24 hours |
| Moderate | 30–50   | 500 to 1500  | Every 24 hours |
| Severe   | <30   | 500 to 1000  | Every 24 hours |

### 3 DOSAGE FORMS AND STRENGTHS

Levetiracetam extended-release tablets, USP are white, modified oval-shaped, convex, film-coated, debossed "4186" on one side and debossed "V" on the reverse side and contain 500 mg levetiracetam.

Levetiracetam extended-release tablets, USP are white, modified oval-shaped, convex, film-coated, debossed "4187" on one side and debossed "V" on the reverse side and contain 750 mg levetiracetam.

### 4 CONTRAINDICATIONS

None.

### 5 WARNINGS AND PRECAUTIONS

#### 5.1 Behavioral Abnormalities and Psychotic Symptoms

Levetiracetam extended-release tablets may cause behavioral abnormalities and psychotic symptoms. Patients treated with levetiracetam extended-release tablets should be monitored for psychiatric signs and symptoms.

#### Behavioral abnormalities

##### Levetiracetam Extended-Release Tablets

A total of 7% of levetiracetam extended-release tablet-treated patients experienced non-psychotic behavioral disorders (reported as irritability and aggression) compared to 0% of placebo-treated patients. Irritability and aggression was reported in 7% of levetiracetam extended-release tablet-treated patients. Aggression was reported in 1% of levetiracetam extended-release tablet-treated patients.

No patient discontinued treatment or had a dose reduction as a result of these adverse reactions.

The number of patients exposed to levetiracetam extended-release tablets was considerably smaller than the number of patients exposed to immediate-release levetiracetam tablets in controlled trials. Therefore, certain adverse reactions observed in the immediate-release levetiracetam controlled trials will likely occur in patients receiving levetiracetam extended-release tablets.

##### Immediate-Release Levetiracetam Tablets

A total of 13% of adult patients and 38% of pediatric patients (4 to 16 years of age) treated with immediate-release levetiracetam tablets experienced non-psychotic behavioral symptoms (reported as aggression, agitation, anger, anxiety, apathy, depersonalization, depression, emotional lability, hostility, hyperkinesias, irritability, nervousness, neurosis, and personality disorder), compared to 6% and 19% of adult and pediatric patients on placebo. A randomized, double-blind, placebo-controlled study was conducted to assess the neurocognitive and behavioral effects of immediate-release levetiracetam tablets as adjunctive therapy in pediatric patients (4 to 16 years of age). An exploratory analysis suggested a worsening in aggressive behavior in patients treated with immediate-release levetiracetam tablets in that study (See *Use in Specific Populations* (8.4)).

A total of 1.7% of adult patients treated with immediate-release levetiracetam tablets discontinued treatment due to behavioral adverse reactions, compared to 0.2% of placebo-treated patients. The treatment dose was reduced in 0.8% of adult patients treated with immediate-release levetiracetam tablets, compared to 0.5% of placebo-treated patients. Overall, 11% of pediatric patients treated with immediate-release levetiracetam tablets experienced behavioral symptoms associated with discontinuation or dose reduction, compared to 6.2% of placebo-treated pediatric patients.

One percent of adult patients and 2% of pediatric patients (4 to 16 years of age) treated with immediate-release levetiracetam tablets experienced psychotic symptoms, compared to 0.2% and 2%, respectively, in adult and placebo-treated pediatric patients. In a controlled study that assessed the neurocognitive and behavioral effects of immediate-release levetiracetam tablets in pediatric patients 4 to 16 years of age, 1.6% Levetiracetam-treated patients experienced paranoia, compared to

## WARNINGS AND PRECAUTIONS

Behavioral abnormalities including psychotic symptoms, suicidal ideation, irritability, and aggressive behavior have been observed; monitor patients for psychiatric signs and symptoms (5.1)

Suicidal Behavior and Ideation: Monitor patients for new or worsening depression, suicidal thoughts/behavior, and/or unusual changes in mood or behavior (5.2)

Monitor for somnolence and fatigue and advise patients not to drive or operate machinery until they have gained sufficient experience on levetiracetam extended-release tablets (5.3)

Withdrawal Seizures: Levetiracetam extended-release tablets must be gradually withdrawn (5.6)

## ADVERSE REACTIONS

Most common adverse reactions (incidence ≥5% more than placebo) include: somnolence and irritability (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Qualified Pharmaceuticals at 1-800-444-0011 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).

## USE IN SPECIFIC POPULATIONS

Pregnancy: Plasma levels of levetiracetam may be decreased and therefore need to be monitored closely during pregnancy. Based on animal data, may cause fetal harm (5.8, 8.1)

## See 17 for PATIENT COUNSELING INFORMATION and MEDICATION GUIDE

Revised: 05/2016

|      |   |
|------|---|
| 8.5  | Geriatric Use   |
| 8.6  | Renal Impairment  |
| 10   | OVERDOSAGE  |
| 10.1 | Signs, Symptoms and Laboratory Findings of Acute Overdosage in Humans |
| 10.2 | Management of Overdose  |
| 10.3 | Hemodialysis  |
| 11   | DESCRIPTION   |
| 12   | CLINICAL PHARMACOLOGY   |
| 12.1 | Mechanism of Action   |
| 12.2 | Pharmacodynamics  |
| 12.3 | Pharmacokinetics  |
| 13   | NONCLINICAL TOXICOLOGY  |
| 13.1 | Carcinogenesis, Mutagenesis, Impairment of Fertility                  |
| 14   | CLINICAL STUDIES  |
| 14.1 | Levetiracetam Extended-Release Tablets in Adults                      |
| 14.2 | Immediate-Release Levetiracetam Tablets in Adults                     |
| 14.3 | Immediate-Release Levetiracetam Tablets in Pediatric Patients         |
| 16   | HOW SUPPLIED/STORAGE AND HANDLING                                     |
| 16.1 | How Supplied  |
| 16.2 | Storage   |
| 17   | PATIENT COUNSELING INFORMATION  |

\*Sections or subsections omitted from the Full Prescribing Information are not listed.

no placebo-treated patients. There were 3.1% patients treated with immediate-release levetiracetam tablets who experienced confusional state, compared to no placebo-treated patients (See *Use in Specific Populations* (8.4)).

#### Psychotic symptoms

##### Immediate-Release Levetiracetam Tablets

One percent of levetiracetam-treated adult patients experienced psychotic symptoms compared to 0.2% of placebo-treated patients.

Two (0.3%) immediate-release levetiracetam tablets-treated adult patients were hospitalized and their treatment was discontinued due to psychosis. Both events, reported as psychosis, developed within the first week of treatment and resolved within 1 to 2 weeks following treatment discontinuation. There was no difference between drug and placebo-treated patients in the incidence of pediatric patients who discontinued treatment due to psychotic and non-psychotic adverse reactions.

##### 5.2 Suicidal Behavior and Ideation

Antiepileptic drugs (AEDs), including levetiracetam extended-release tablets, increase the risk of suicidal thoughts or behavior in patients taking these drugs for any indication. Patients treated with any AED for any indication should be monitored for the emergence or worsening of depression, suicidal thoughts or behavior, and/or any unusual changes in mood or behavior.

Pool analyses of 149 placebo-controlled clinical trials (mono- and adjunctive therapy) of 11 different AEDs showed that patients randomized to one of the AEDs had approximately twice the risk (adjusted Relative Risk 1.8, 95% CI: 1.2, 2.7) of suicidal thinking or behavior compared to patients randomized to placebo. In these trials, which had a median treatment duration of 12 weeks, the estimated incidence rate of suicidal behavior or ideation among 27,863 AED-treated patients was 0.43%, compared to 0.24% among 16,029 placebo-treated patients, representing an increase of approximately one case of suicidal thinking or behavior for every 530 patients treated. There were four suicides in drug-treated patients in the trials and none in placebo-treated patients, but the number is too small to allow any conclusion about drug effect on suicide.

The increased risk of suicidal thoughts or behavior with AEDs was observed as early as one week after starting drug treatment with AEDs and persisted for the duration of treatment assessed. Because most trials included in the analysis did not extend beyond 24 weeks, the risk of suicidal thoughts or behavior beyond 24 weeks could not be assessed.

The risk of suicidal thoughts or behavior was generally consistent among drugs in the data analyzed. The finding of increased risk with AEDs of varying mechanisms of action and across a range of indications suggests that the risk applies to all AEDs used for any indication. The risk did not vary substantially by age (5 to 100 years) in the clinical trials analyzed. Table 2 shows absolute and relative risk by indication for all evaluated AEDs.

Table 2. Risk by Indication for Antiepileptic Drugs in the Pooled Analysis

| Indication  | Placebo Patients with Events Per 1000 Patients | Drug Patients with Events Per 1000 Patients | Relative Risk: Incidence of Events in Drug Patients/Incidence in Placebo Patients | Risk Difference: Additional Drug Patients with Events Per 1000 Patients |
|-------------|--|---|---|---|
| Epilepsy    | 1.0  | 3.4   | 3.5   | 2.4   |
| Psychiatric | 5.7  | 8.5   | 1.5   | 2.9   |
| Other       | 1.0  | 1.8   | 1.9   | 0.9   |
| Total       | 2.4  | 4.3   | 1.8   | 1.9   |

The relative risk for suicidal thoughts or behavior was higher in clinical trials for epilepsy than in clinical trials for psychiatric or other conditions, but the absolute risk differences were similar for the epilepsy and psychiatric indications.

Anyone considering prescribing levetiracetam extended-release tablets or any other AED must balance the risk of suicidal thoughts or behavior with the risk of untreated illness. Epilepsy and many other illnesses for which AEDs are prescribed are themselves associated with morbidity and mortality and an increased risk of suicidal thoughts and behavior. Should suicidal thoughts and behavior emerge during treatment, the prescriber needs to consider whether the emergence of these symptoms in any given patient may be related to the illness being treated.

Patients, their caregivers, and families should be informed that AEDs increase the risk of suicidal thoughts and behavior and should be advised of the need to be alert for the emergence or worsening of the signs and symptoms of depression, any unusual changes in mood or behavior, or the emergence of suicidal thoughts, behavior, or thoughts about self-harm. Behaviors of concern should be reported immediately to healthcare providers.

##### 5.3 Somnolence and Fatigue

Levetiracetam extended-release tablets may cause somnolence and fatigue. Patients should be monitored for these signs and symptoms and advised not to drive or operate machinery until they have gained sufficient experience on levetiracetam extended-release tablets to gauge whether it adversely affects their ability to drive or operate machinery.

Somnolence

##### Levetiracetam Extended-Release Tablets

In a levetiracetam extended-release tablets double-blind, controlled trial in patients experiencing partial onset seizures, 8% of levetiracetam extended-release tablets-treated patients experienced somnolence compared to 3% of placebo-treated patients.

No patient discontinued treatment or had a dose reduction as a result of these adverse reactions.

The number of patients exposed to levetiracetam extended-release tablets was considerably smaller than the number of patients exposed to immediate-release levetiracetam tablets in controlled trials. Therefore, certain adverse reactions observed in the immediate-release levetiracetam controlled trials will likely occur in patients receiving levetiracetam extended-release tablets.

##### Immediate-Release Levetiracetam Tablets

In controlled trials of adult patients with epilepsy experiencing partial onset seizures, 15% of immediate-release levetiracetam tablet-treated patients reported somnolence, compared to 8% of placebo-treated patients. There was no clear dose response up to 3000 mg/day. In a study where there was no titration, about 45% of patients receiving 4000 mg/day reported somnolence. The somnolence was considered serious in 0.3% of the immediate-release levetiracetam tablets-treated patients, compared to 0% in the placebo group. About 3% of immediate-release levetiracetam tablets-treated patients discontinued treatment due to somnolence, compared to 0.7% of placebo-treated patients. In 1.4% of immediate-release levetiracetam

tablets-treated patients and in 0.9% of placebo-treated patients the dose was reduced, while 0.3% of the treated patients were hospitalized due to somnolence.

#### Asthenia

##### Immediate-Release Levetiracetam Tablets

In controlled trials of adult patients with epilepsy experiencing partial onset seizures, 15% of immediate-release levetiracetam tablets-treated patients reported asthenia, compared to 9% of placebo-treated patients. Treatment was discontinued due to asthenia in 0.8% of immediate-release levetiracetam tablets-treated patients as compared to 0.5% of placebo-treated patients. In 0.5% of immediate-release levetiracetam tablet-treated patients and in 0.2% of placebo-treated patients, the dose was reduced due to asthenia.

Somnolence and asthenia occurred most frequently within the first 4 weeks of treatment.

#### 5.4 Serious Dermatological Reactions

Serious dermatological reactions, including Stevens-Johnson syndrome (SJS) and toxic epidermal necrolysis (TEN), have been reported in patients treated with levetiracetam. The median time of onset is reported to be 14 to 17 days, but cases have been reported at least four months after initiation of treatment. Recurrence of the serious skin reactions following rechallenge with levetiracetam has also been reported. Levetiracetam extended-release tablets should be discontinued at the first sign of a rash, unless the rash is clearly not drug-related. If signs or symptoms suggest SJS/TEN, use of this drug should not be resumed and alternative therapy should be considered.

#### 5.5 Coordination Difficulties

Coordination difficulties were not observed in a levetiracetam extended-release tablets controlled trial, however, the number of patients exposed to levetiracetam extended-release tablets was considerably smaller than the number of patients exposed to immediate-release levetiracetam tablets in controlled trials. However, adverse reactions observed in the immediate-release levetiracetam controlled trials may also occur in patients receiving levetiracetam extended-release tablets.

#### Immediate-Release Levetiracetam Tablets

A total of 3.4% of adult levetiracetam-treated patients experienced coordination difficulties, (reported as either ataxia, abnormal gait, or incoordination) compared to 1.6% of placebo-treated patients. A total of 0.4% of patients in controlled trials discontinued levetiracetam treatment due to ataxia, compared to 0% of placebo-treated patients. In 0.7% of levetiracetam-treated patients and in 0.2% of placebo-treated patients, the dose was reduced due to coordination difficulties, while one of the levetiracetam-treated patients was hospitalized due to worsening of pre-existing ataxia. These events occurred most frequently within the first 4 weeks of treatment.

Patients should be monitored for these signs and symptoms and advised not to drive or operate machinery until they have gained sufficient experience on levetiracetam to gauge whether it could adversely affect their ability to drive or operate machinery.

#### 5.6 Withdrawal Seizures

Antiepileptic drugs, including levetiracetam extended-release tablets, should be withdrawn gradually to minimize the potential of increased seizure frequency.

#### 5.7 Hematologic Abnormalities

Levetiracetam extended-release tablets can cause hematologic abnormalities. Hematologic abnormalities occurred in clinical trials and included decreases in red blood cell (RBC) counts, hemoglobin, and hematocrit, and increases in eosinophil counts. Decreased white blood cell (WBC) and neutrophil counts also occurred in clinical trials. Cases of agranulocytosis have been reported in the postmarketing setting.

In controlled trials of immediate-release levetiracetam tablets in patients experiencing partial onset seizures, minor, but statistically significant, decreases compared to placebo in total mean RBC count (0.03 × 10<sup>12</sup>/mm<sup>3</sup>), mean hemoglobin (0.09 g/dL), and mean hematocrit (0.38%), were seen in immediate-release levetiracetam-treated patients.

A total of 3.2% of levetiracetam-treated and 1.8% of placebo-treated patients had at least one possibly significant (<2.8 × 10<sup>12</sup>/L decreased WBC, and 2.4% of levetiracetam-treated and 1.4% of placebo-treated patients had at least one possibly significant (<1.0 × 10<sup>9</sup>/L decreased neutrophil count). Of the levetiracetam-treated patients with a low neutrophil count, all but one rose towards or baseline with continued treatment. No patient was discontinued secondary to low neutrophil counts.

In pediatric patients (4 to <16 years of age), statistically significant decreases in WBC and neutrophil counts were seen in patients treated with immediate-release levetiracetam tablets, as compared to placebo. The mean decreases from baseline in the immediate-release levetiracetam tablets group were -0.4 × 10<sup>9</sup>/L and -0.3 × 10<sup>9</sup>/L, respectively, whereas there were small increases in the placebo group. A significant increase in mean relative lymphocyte counts was observed in 1.7% of patients treated with immediate-release levetiracetam tablets compared to a decrease of 4% in patients on placebo.

In a controlled pediatric trial, a possibly clinically significant abnormal low WBC value was observed in 3% of patients treated with immediate-release levetiracetam tablets, compared to no patients on placebo. However, there was no apparent difference between treatment groups with respect to neutrophils. No patient was discontinued secondary to low WBC or neutrophil counts.

In a controlled pediatric cognitive and neuropsychological safety study, two subjects (6.1%) in the placebo group and 5 subjects (8.6%) in the immediate-release levetiracetam tablets-treated group had high eosinophil count values that were possibly clinically significant (≥10% or ≥0.7X10<sup>9</sup>/L).

#### 5.8 Seizure Control During Pregnancy

Physiological changes may gradually decrease plasma levels of levetiracetam throughout pregnancy. This decrease is more pronounced during the third trimester. It is recommended that patients be monitored carefully during pregnancy. Close monitoring should continue through the postpartum period especially if the dose was changed during pregnancy.

## 6 ADVERSE REACTIONS

The following adverse reactions are discussed in more details in other sections of labeling:

- Behavioral abnormalities and Psychotic Symptoms (See *Warnings and Precautions* (5.1))
- Suicidal Behavior And Ideation (See *Warnings and Precautions* (5.2))
- Somnolence And Fatigue (See *Warnings and Precautions* (5.3))
- Serious Dermatological Reactions (See *Warnings and Precautions* (5.4))
- Coordination Difficulties (See *Warnings and Precautions* (5.5))
- Hematologic Abnormalities (See *Warnings and Precautions* (5.7))

#### 6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug can be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

#### Levetiracetam Extended-Release Tablets

In a controlled clinical trial in patients with partial onset seizures, the most common adverse reactions in patients receiving levetiracetam extended-release tablets in combination with other AEDs, for events with rates greater than placebo, were irritability and somnolence.

Table 3 lists adverse reactions that occurred in at least 5% of epilepsy patients receiving levetiracetam extended-release tablets in a placebo-controlled study and were numerically more common than in patients treated with placebo. In this study, either levetiracetam extended-release tablets or placebo were added to concurrent AED therapy.

Table 3. Adverse Reactions in A Placebo-Controlled, Add-On Study In Patients Experiencing Partial Onset Seizures

|                 | Levetiracetam Extended-Release Tablets (N=77) % | Placebo (N=79) % |
|-----------------|---|------------------|
| Influenza       | 8   | 4                |
| Somnolence      | 8   | 3                |
| Irritability    | 7   | 0                |
| Nasopharyngitis | 7   | 5                |
| Dizziness       | 5   | 3                |
| Nausea          | 5   | 3                |

#### Discontinuation or Dose Reduction in a Levetiracetam Extended-Release Tablets Controlled Clinical Study

In a controlled clinical study, 5% of patients receiving levetiracetam extended-release tablets and 3% receiving placebo discontinued as a result of an adverse reaction. The adverse reactions that resulted in discontinuation and that occurred more frequently in levetiracetam extended-release tablet-treated patients than in placebo-treated patients were asthenia, epilepsy, seizure table-treated patient and no placebo-treated patients.

#### Immediate-Release Levetiracetam Tablets

Table 4 lists the adverse reactions in controlled studies of immediate-release levetiracetam tablets in adult patients experiencing partial onset seizures. Although the pattern of adverse reactions in a levetiracetam extended-release tablets study seems somewhat different from that seen in partial onset seizure controlled studies for immediate-release levetiracetam tablets, this is possibly due to the much smaller number of patients in this study compared to the immediate-release tablet studies. The adverse reactions for mouth ulceration, rash and respiratory failure. Each of these adverse reactions led to discontinuation in a levetiracetam extended-release tablet-treated patient and no placebo-treated patients.

Table 4 lists adverse reactions that occurred in at least 1% of adult epilepsy patients receiving immediate-release levetiracetam tablets in placebo-controlled studies and were numerically more common than in patients treated with placebo. In these studies, either immediate-release levetiracetam tablets or placebo were added to concurrent AED therapy.

Table 4. Adverse Reactions in Pooled Placebo-Controlled, Add-On Studies in Adults Experiencing Partial Onset Seizures

|                    | Immediate-Release Levetiracetam Tablets (N=789) % | Placebo (N=439) % |
|--------------------|---|-------------------|
| Asthenia           | 15  | 9                 |
| Somnolence         | 15  | 8                 |
| Headache           | 14  | 13                |
| Infection          | 13  | 8                 |
| Dizziness          | 9   | 4                 |
| Pain               | 7   | 6                 |
| Pharyngitis        | 6   | 4                 |
| Depression         | 4   | 2                 |
| Nervousness        | 4   | 2                 |
| Rhinitis           | 4   | 3                 |
| Anorexia           | 3   | 2                 |
| Alzaco             | 3   | 1                 |
| Vertigo            | 3   | 1                 |
| Amnesia            | 2   | 1                 |
| Anxiety            | 2   | 1                 |
| Cough Increased    | 2   | 1                 |
| Diplopia           | 2   | 1                 |
| Emotional Lability | 2   | 0                 |
| Hostility          | 2   | 1                 |
| Paresthesia        | 2   | 1                 |
| Sinusitis          | 2   | 1                 |

#### Pediatric Patients 4 Years to <16 Years

In a pooled analysis of two controlled pediatric clinical studies in children 4 to 16 years of age with partial onset seizures, the adverse reactions most frequently reported with the use of immediate-release levetiracetam tablets in combination with other AEDs, and with greater frequency than in patients on placebo, were fatigue, aggression, nasal congestion, decreased appetite, and irritability.

Table 5 lists adverse reactions that occurred in at least 2% of pediatric patients treated with immediate-release levetiracetam tablets and were more common than in pediatric patients on placebo. In these studies, either immediate-release levetiracetam tablets or placebo was added to concurrent AED therapy. Adverse reactions were usually mild to moderate in intensity.

Table 5. Adverse Reactions in Pooled Placebo-Controlled, Add-On Studies in Pediatric Patients Ages 4 to 16 Years Experiencing Partial Onset Seizures

|                 | Immediate-Release Levetiracetam Tablets (N=165) % | Placebo (N=131) % |
|-----------------|---|-------------------|
| Headache        | 19  | 15                |
| Nasopharyngitis | 15  | 12                |
| Vomiting        | 15  | 12                |
| Somnolence      | 13  | 9                 |
| Fatigue         | 11  | 5                 |
| Aggression      |   |                   |

related analyses showed a rank order of affinity for SV2A which correlated with the potency of their antiseizure activity in audiogenic seizure-prone mice. These findings suggest that the interaction of levetiracetam with the SV2A protein may contribute to the antiepileptic mechanism of action of the drug.

## 12.2 Pharmacodynamics

### Effects on QTc Interval

The effects of levetiracetam extended-release tablets on QTc prolongation is expected to be the same as that of immediate-release levetiracetam. The effect of immediate-release levetiracetam on QTc prolongation was evaluated in a randomized, double-blind, placebo-controlled (moxifloxacin 400 mg) and placebo-controlled crossover study of immediate-release levetiracetam (1000 mg or 5000 mg) in 52 healthy subjects. The upper bound of the 90% confidence interval for the largest placebo-adjusted, baseline-corrected QTc was below 10 milliseconds. Therefore, there was no evidence of significant QTc prolongation in this study.

## 12.3 Pharmacokinetics

### Overview

Bioavailability of levetiracetam extended-release tablets is similar to that of the immediate-release levetiracetam tablets. The pharmacokinetics (AUC and  $C_{max}$ ) were shown to be dose proportional after single dose administration of 1000 mg, 2000 mg, and 3000 mg extended-release levetiracetam. Plasma half-life of extended-release levetiracetam is approximately 7 hours.

Levetiracetam is almost completely absorbed after oral administration. The pharmacokinetics of levetiracetam are linear and time-invariant, with low intra- and inter-subject variability. Levetiracetam is not significantly protein-bound (<10% bound) and its volume of distribution is close to the volume of intracellular and extracellular water. Sixty-six percent (66%) of the dose is renally excreted unchanged. The major metabolic pathway of levetiracetam (24% of dose) is an enzymatic hydrolysis of the acetamide group. It is not liver cytochrome P450 dependent. The metabolites have no known pharmacological activity and are renally excreted. Plasma half-life of levetiracetam across studies is approximately 6 to 8 hours. The half-life is increased in the elderly (primarily due to impaired renal clearance) and in subjects with renal impairment.

### Absorption and Distribution

Extended-release levetiracetam peak plasma concentrations occur in about 4 hours. The time to peak plasma concentrations is about 3 hours longer with extended-release levetiracetam than with immediate-release tablets.

Single administration of two 500 mg extended-release levetiracetam tablets once daily produced comparable maximal plasma concentrations and area under the plasma concentration versus time as did the administration of one 500 mg immediate-release tablet twice daily in fasting conditions. After multiple dose extended-release levetiracetam tablets intake, extent of exposure (AUC<sub>0-∞</sub>) was similar to extent of exposure after multiple dose immediate-release tablets intake.  $C_{max}$  and  $C_{min}$  were lower by 17% and 26% after multiple dose extended-release levetiracetam tablets intake in comparison to multiple dose immediate-release tablets intake. Intake of a high fat, high calorie breakfast before the administration of extended-release levetiracetam tablets resulted in a higher peak concentration, and longer median time to peak. The median time to peak ( $T_{max}$ ) was 2 hours longer in the fed state.

Two 750 mg extended-release levetiracetam tablets were bioequivalent to a single administration of three 500 mg extended-release levetiracetam tablets.

### Metabolism

Levetiracetam is not extensively metabolized in humans. The major metabolic pathway is the enzymatic hydrolysis of the acetamide group, which produces the carboxylic acid metabolite, uch L057 (24% of dose) and is not dependent on any liver cytochrome P450 isoenzymes. The major metabolite is inactive in animal seizure models. Two minor metabolites were identified as the product of hydrolyzation of the 2-oxo-pyrrolidine ring (2% of dose and opening of the 2-oxo-pyrrolidine ring in position 5 (1% of dose). There is no enantiomeric interconversion of levetiracetam or of its major metabolite.

### Elimination

Levetiracetam plasma half-life in adults is 7 ± 1 hour and is unaffected by either dose or repeated administration. Levetiracetam is eliminated from the systemic circulation by renal excretion as unchanged drug which represents 86% of administered dose. The total body clearance is 0.96 mL/min/kg and the renal clearance is 0.6 mL/min/kg. The mechanism of excretion is glomerular filtration with subsequent partial tubular reabsorption. The metabolite uch L057 is excreted by glomerular filtration and active tubular secretion with a renal clearance of 4 mL/min/kg. Levetiracetam elimination is correlated to creatinine clearance. Levetiracetam clearance is reduced in patients with impaired renal function (see *Dosage and Administration (2.2) and Use in Specific Populations (8.5)*).

### Specific Populations

#### Elderly

There are insufficient pharmacokinetic data to specifically address the use of extended-release levetiracetam in the elderly population.

Pharmacokinetics of immediate-release levetiracetam were evaluated in 16 elderly subjects (age 61 to 88 years) with creatinine clearance ranging from 30 to 74 mL/min. Following oral administration of twice-daily dosing for 10 days, total body clearance decreased by 38% and the half-life was 2.5 hours longer in the elderly compared to healthy adults. This is most likely due to the decrease in renal function in these subjects.

#### Pediatric Patients

An open label, multicenter, parallel-group, two-arm study was conducted to evaluate the pharmacokinetics of levetiracetam extended-release tablets in pediatric patients (13 to 16 years old) and in adults (18 to 55 years old) with epilepsy. Levetiracetam extended-release oral tablets (1000 mg to 3000 mg) were administered once daily with a minimum of 4 days and a maximum of 7 days of treatment to 12 pediatric patients and 12 adult patients. Study Dose-normalized steady-state exposure parameters,  $C_{max}$  and AUC, were comparable between pediatric and adult patients.

#### Pregnancy

Levetiracetam levels may decrease during pregnancy.

#### Gender

Extended-release levetiracetam  $C_{max}$  was 21 to 30% higher and AUC was 8 to 18% higher in women (N=12) compared to men (N=12). However, clearances adjusted for body weight were comparable.

#### Race

Formal pharmacokinetic studies of the effects of race have not been conducted with extended-release or immediate-release levetiracetam. Cross study comparisons involving Caucasians (N=12) and Asians (N=12), however, show that pharmacokinetics of immediate-release levetiracetam were comparable between the two races. Because levetiracetam is primarily renally excreted and there are no important racial differences in creatinine clearance, pharmacokinetic differences due to race are not expected.

#### Renal Impairment

The effect of levetiracetam extended-release tablets on renally impaired patients was not assessed in a controlled study. However, it is expected that the effect on levetiracetam extended-release tablet-treated patients would be similar to that seen in controlled studies of immediate-release levetiracetam tablets. In patients with end stage renal disease on dialysis, it is recommended that immediate-release levetiracetam be used instead of levetiracetam extended-release tablets.

The disposition of immediate-release levetiracetam was studied in adult subjects with varying degrees of renal function. Total body clearance of levetiracetam is reduced in patients with impaired renal function by 40% in the mild group (CL<sub>cr</sub> = 30 to 80 mL/min), 50% in the moderate group (CL<sub>cr</sub> = 30 to 50 mL/min) and 60% in the severe renal impairment group (CL<sub>cr</sub> <30 mL/min). Clearance of levetiracetam is correlated with creatinine clearance.

In anuric (end stage renal disease) patients, the total body clearance decreased 70% compared to normal subjects (CL<sub>cr</sub> >80 mL/min). Approximately 50% of the pool of levetiracetam in the body is removed during a standard 4-hour hemodialysis procedure (see *Dosage and Administration (2.2)*).

#### Hepatic Impairment

In subjects with mild (Child-Pugh A) to moderate (Child-Pugh B) hepatic impairment, the pharmacokinetics of levetiracetam were unchanged. In patients with severe hepatic impairment (Child-Pugh C), total body clearance was 50% that of normal subjects, but decreased renal clearance accounted for most of the decrease. No dose adjustment is needed for patients with hepatic impairment.

#### Drug Interactions

*In vitro* data on metabolic interactions indicate that levetiracetam is unlikely to produce, or be subject to, pharmacokinetic interactions. Levetiracetam and its major metabolite, at concentrations well above  $C_{max}$  levels achieved within the therapeutic dose range, are neither inhibitors nor, nor high affinity substrates for, human liver cytochrome P450 isoforms, epoxide hydrolase or UDP-glucuronation enzymes. In addition, levetiracetam does not affect the *in vitro* glucuronation of valproic acid.

Potential pharmacokinetic interactions of or with levetiracetam were assessed in clinical pharmacokinetic studies (phenytoin, valproate, warfarin, digoxin, oral contraceptive, probenecid) and through pharmacokinetic screening with immediate-release levetiracetam tablets in a placebo-controlled clinical studies in epilepsy patients. The potential for drug interactions for levetiracetam extended-release tablets is expected to be essentially the same as that with immediate-release levetiracetam tablets.

#### Phenytoin

Immediate-release levetiracetam tablets (3000 mg daily) had no effect on the pharmacokinetic disposition of phenytoin in patients with refractory epilepsy. Pharmacokinetics of levetiracetam were also not affected by phenytoin.

#### Valproate

Immediate-release levetiracetam tablets (1500 mg twice daily) did not alter the pharmacokinetics of valproate in healthy volunteers. Valproate 500 mg twice daily did not modify the rate or extent of levetiracetam absorption or its plasma clearance or urinary excretion. There was also no effect on exposure to and the excretion of the primary metabolite, uch L057.

#### Other Antiepileptic Drugs

Potential drug interactions between immediate-release levetiracetam tablets and other AEDs (carbamazepine, gabapentin, lamotrigine, phenobarbital, phenytoin, primidone and valproate) were also assessed by evaluating the serum concentrations of levetiracetam and these AEDs during placebo-controlled clinical studies. These data indicate that levetiracetam does not influence the plasma concentration of other AEDs and that these AEDs do not influence the pharmacokinetics of levetiracetam.

#### Oral Contraceptives

Immediate-release levetiracetam tablets (500 mg twice daily) did not influence the pharmacokinetics of an oral contraceptive containing 0.03 mg ethinyl estradiol and 0.15 mg levonorgestrel, or of the luteinizing hormone and progesterone levels, indicating that impairment of contraceptive efficacy is unlikely. Coadministration of this oral contraceptive did not influence the pharmacokinetics of levetiracetam.

#### Digoxin

Immediate-release levetiracetam tablets (1000 mg twice daily) did not influence the pharmacokinetics and pharmacodynamics (EC<sub>50</sub>) of digoxin given as a 0.25 mg dose every day. Coadministration of digoxin did not influence the pharmacokinetics of levetiracetam.

#### Warfarin

Immediate-release levetiracetam tablets (1000 mg twice daily) did not influence the pharmacokinetics of R and S warfarin. Prothrombin time was not affected by levetiracetam. Coadministration of warfarin did not affect the pharmacokinetics of levetiracetam.

#### Probenecid

Probenecid, a renal tubular secretion blocking agent, administered at a dose of 500 mg four times a day, did not change the pharmacokinetics of levetiracetam 1000 mg twice daily.  $C^*_{max}$  of the metabolite, uch L057, was approximately doubled in the presence of probenecid while the fraction of drug excreted unchanged in the urine remained the same. Renal clearance of uch L057 in the presence of probenecid decreased 60%, probably related to competitive inhibition of tubular secretion of uch L057. The effect of immediate-release levetiracetam tablets on probenecid was not studied.

## 13 NONCLINICAL TOXICOLOGY

### 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

#### Carcinogenesis

Rats were dosed with levetiracetam in the diet for 104 weeks at doses of 50, 300 and 1800 mg/kg/day. The highest dose is 6 times the maximum recommended daily human dose (MRHD) of 3000 mg on a mg/m<sup>2</sup> basis and it also provided systemic exposure (AUC) approximately 6 times that achieved in humans receiving the MRHD. There was no evidence of carcinogenicity. In mice, oral administration of levetiracetam for 80 weeks (doses up to 960 mg/kg/day) or 2 years (doses up to 4000 mg/kg/day, lowered to 3000 mg/kg/day after 45 weeks due to intolerability) was not associated with an increase in tumors. The highest dose tested in mice for 2 years (3000 mg/kg/day) is approximately 5 times the MRHD on a mg/m<sup>2</sup> basis.

#### Mutagenesis

Levetiracetam was not mutagenic in the Ames test or in mammalian cells *in vitro* in the Chinese hamster ovary/HCPRT locus assay. It was not clastogenic in an *in vitro* analysis of metaphase chromosomes obtained from Chinese hamster ovary cells or in an *in vivo* micronucleus assay. The hydrolysis product and major human metabolite of levetiracetam (uch L057) was not mutagenic in the Ames test or in the *in vitro* mouse lymphoma assay.

#### Impairment of Fertility

No adverse effects on male or female fertility or reproductive performance were observed in rats at oral doses up to 1800 mg/kg/day (6 times the maximum recommended human dose on a mg/m<sup>2</sup> systemic exposure (AUC) basis).

## 14 CLINICAL STUDIES

The effectiveness of levetiracetam extended-release tablets as adjunctive therapy in partial onset seizures in adults was established in one multicenter, randomized, double-blind, placebo-controlled clinical study in patients who had refractory partial onset seizures with or without secondary generalization. This was supported by the demonstration of efficacy of immediate-release levetiracetam tablets (see below) in partial seizures in three multicenter, randomized, double-blind, placebo-controlled clinical studies in adults, as well as a demonstration of comparable bioavailability between the extended-release and immediate-release formulations (see *Clinical Pharmacology (12.3)*) in adults. The effectiveness for levetiracetam extended-release tablets as adjunctive therapy in partial onset seizures in pediatric patients, 12 years of age and older, was based upon a single pharmacokinetic study showing comparable pharmacokinetics of levetiracetam extended-release tablets in adults and adolescents (see *Clinical Pharmacology (12.3)*). All studies are described below.

### 14.1 Levetiracetam Extended-Release Tablets in Adults

The effectiveness of levetiracetam extended-release tablets as adjunctive therapy (added to other antiepileptic drugs) was established in one multicenter, randomized, double-blind, placebo-controlled clinical study across 7 countries in patients who had refractory partial onset seizures with or without secondary generalization (Study 1).

#### Study 1

Patients enrolled in Study 1 had at least eight partial seizures with or without secondary generalization during the 8-week baseline period and at least two partial seizures in each 4-week interval of the baseline period. Patients were taking a stable dose regimen of at least one AED, and could take a maximum of three AEDs. After a prospective baseline period of 8 weeks, 158 patients were randomized to placebo (N=79) or 1000 mg (two 500 mg tablets) of levetiracetam extended-release tablets (N=79), given once daily over a 12-week treatment period.

The primary efficacy endpoint in Study 1 was the percent reduction over placebo in mean weekly frequency of partial onset seizures. The median percent reduction in weekly partial onset seizure frequency from baseline over the treatment period was 46.1% in the levetiracetam extended-release tablets 1000 mg treatment group (N=79) and 33.4% in the placebo group (N=78). The estimated percent reduction over placebo in weekly partial onset seizure frequency over the treatment period was 14.4% (statistically significant).

The relationship between the effectiveness of the same daily dose of levetiracetam extended-release tablets and immediate-release levetiracetam tablets has not been studied and is unknown.

### 14.2 Immediate-Release Levetiracetam Tablets in Adults

The effectiveness of immediate-release levetiracetam tablets as adjunctive therapy (added to other antiepileptic drugs) in adults was established in three multicenter, randomized, double-blind, placebo-controlled clinical studies in patients who had refractory partial onset seizures with or without secondary generalization (Studies 2, 3, and 4). The tablet formulation was used in all three studies. In these studies, 504 patients were randomized to placebo, levetiracetam tablets 1000 mg, levetiracetam tablets 2000 mg, or levetiracetam tablets 3000 mg/day. Patients enrolled in Study 2 or Study 3 had refractory partial onset seizures for at least two years, and had taken two or more AEDs. Patients enrolled in Study 4 had refractory partial onset seizures for at least 1 year and had taken one AED. At the time of the study, patients were taking a stable dose regimen of at least one AED, and could take a maximum of two AEDs. During the baseline period, patients had to have experienced at least two partial onset seizures during each 4-week period.

#### Study 2

Study 2 was a double-blind, placebo-controlled, parallel-group study conducted at 41 sites in the United States, comparing immediate-release levetiracetam tablets 1000 mg/day (N=97), immediate-release levetiracetam tablets 3000 mg/day (N=101), and placebo (N=95), given in equally divided doses twice daily. After a prospective baseline period of 12 weeks, patients in Study 2 were randomized to one of the three treatment groups described above. The 18-week treatment period consisted of a 6-week titration period, followed by a 12-week fixed dose evaluation period, during which concomitant AED regimens were held constant. The primary measure of effectiveness in Study 2 was a between-group comparison of the percent reduction in weekly partial seizure frequency relative to placebo over the entire randomized treatment period (titration + evaluation period). Secondary outcome variables included the responder rate (incidence of patients with ≥50% reduction from baseline in partial onset seizure frequency). The results of Study 2 are displayed in Table 6.


Table 6: Reduction in Mean Over Placebo in Weekly Frequency Of Partial Onset Seizures in Study 2

|   | Placebo (N=95) | Immediate-Release Levetiracetam Tablets 1000 mg/day (N=97) | Immediate-Release Levetiracetam Tablets 3000 mg/day (N=101) |
|---|----------------|--|---|
| Percent reduction in partial seizure frequency over placebo | –              | 26.1%*   | 30.1%*  |

\*statistically significant versus placebo

The percentage of patients (y-axis) who achieved ≥50% reduction from baseline in weekly partial onset seizure frequency over the entire randomized treatment period (titration + evaluation period) within the three treatment groups (x-axis) in Study 2 is presented in Figure 1.

Figure 1: Responder Rate (≥50% Reduction From Baseline) in Study 2



\* statistically significant versus placebo

#### Study 3

Study 3 was a double-blind, placebo-controlled, crossover study conducted at 62 centers in Europe, comparing immediate-release levetiracetam tablets 1000 mg/day (N=106), immediate-release levetiracetam tablets 2000 mg/day (N=105), and placebo (N=111), given in equally divided doses twice daily.

The first period of the study (Period A) was designed to be analyzed as a parallel-group study. After a prospective baseline period of up to 12 weeks, patients in Study 3 were randomized to one of the three treatment groups described above. The 16-week treatment period consisted of the 4-week titration period followed by a 12-week fixed dose evaluation period, during which concomitant AED regimens were held constant. The primary measure of effectiveness in Study 3 was a between-group comparison of the percent reduction in weekly partial seizure frequency relative to placebo over the entire randomized treatment period (titration + evaluation period). Secondary outcome variables included the responder rate (incidence of patients with ≥50% reduction from baseline in partial onset seizure frequency). The results of the analysis of Period A are displayed in Table 7.


Table 7: Reduction in Mean Over Placebo in Weekly Frequency Of Partial Onset Seizures in Study 3: Period A

|   | Placebo (N=111) | Immediate-Release Levetiracetam Tablets 1000 mg/day (N=106) | Immediate-Release Levetiracetam Tablets 2000 mg/day (N=105) |
|---|-----------------|---|---|
| Percent reduction in partial seizure frequency over placebo | –               | 17.1%*  | 21.4%*  |

\*statistically significant versus placebo

The percentage of patients (y-axis) who achieved ≥50% reduction from baseline in weekly partial onset seizure frequency over the entire randomized treatment period (titration + evaluation period) within the three treatment groups (x-axis) in Study 3 is presented in Figure 2.

Figure 2: Responder Rate (≥50% Reduction From Baseline) in Study 3: Period A



\* statistically significant versus placebo

The comparison of immediate-release levetiracetam tablets 2000 mg/day to immediate-release levetiracetam tablets 1000 mg/day for responder rate in Study 3 was statistically significant (P=0.02). Analysis of the trial as a cross-over study yielded similar results.

#### Study 4

Study 4 was a double-blind, placebo-controlled, parallel-group study conducted at 47 centers in Europe comparing immediate-release levetiracetam tablets 1000 mg treatment group (N=180) and placebo (N=104) in patients with refractory partial onset seizures, with or without secondary generalization, receiving only one concomitant AED. Study drug was given in two divided doses. After a prospective baseline period of 12 weeks, patients in Study 4 were randomized to one of two treatment groups described above. The 16-week treatment period consisted of a 4-week titration period, followed by a 12-week fixed dose evaluation period, during which concomitant AED doses were held constant. The primary measure of effectiveness in Study 4 was a between-group comparison of the percent reduction in weekly seizure frequency relative to placebo over the entire randomized treatment period (titration + evaluation period). Secondary outcome variables included the responder rate (incidence of patients with ≥50% reduction from baseline in partial onset seizure frequency). Table 8 displays the results of Study 4.


Table 8: Reduction in Mean Over Placebo in Weekly Frequency Of Partial Onset Seizures in Study 4

|   | Placebo (N=104) | Immediate-Release Levetiracetam Tablets 3000 mg/day (N=180) |
|---|-----------------|---|
| Percent reduction in partial seizure frequency over placebo | –               | 23.0%*  |

\*statistically significant versus placebo

The percentage of patients (y-axis) who achieved ≥50% reduction from baseline in weekly partial onset seizure frequency over the entire randomized treatment period (titration + evaluation period) within the two treatment groups (x-axis) in Study 4 is presented in Figure 3.

Figure 3: Responder Rate (≥50% Reduction From Baseline) in Study 4



\* statistically significant versus placebo

### 14.3 Immediate-Release Levetiracetam Tablets in Pediatric Patients 4 Years to 16 Years

The use of levetiracetam extended-release tablets in pediatric patients 12 years of age and older is supported by Study 5, which was conducted using immediate-release levetiracetam tablets. Levetiracetam extended-release tablets are not indicated in children below 12 years of age.

#### Study 5

The effectiveness of immediate-release levetiracetam tablets as adjunctive therapy in pediatric patients was established in a multicenter, randomized double-blind, placebo-controlled study, conducted at 60 sites in North America, in children 4 to 16 years of age with partial seizures uncontrolled by standard antiepileptic drugs (Study 5). Eligible patients on a stable dose of 1-2 AEDs, who still experienced at least 4 partial onset seizures during the 4 weeks prior to screening, as well as at least 4 partial onset seizures in each of the two 4-week baseline periods, were randomized to receive either immediate-release levetiracetam tablets or placebo. The enrolled population included 198 patients (levetiracetam tablets N=101; placebo N=97) with refractory partial onset seizures, with or without secondary generalization. Study 5 consisted of an 8-week baseline period and 4-week titration period followed by a 10-week evaluation period. Dosing was initiated at a dose of 20 mg/kg/day in two divided doses. During the treatment period, the immediate-release levetiracetam tablets doses were adjusted in 20 mg/kg/day increments, at 2-week intervals to the target dose of 60 mg/kg/day. The primary measure of effectiveness in Study 5 was a between-group comparison of the percent reduction in weekly partial seizure frequency relative to placebo over the entire 14-week randomized treatment period (titration + evaluation period). Secondary outcome variables included the responder rate (incidence of patients with ≥ 50% reduction from baseline in partial onset seizure frequency per week). Table 9 displays the results of this study.


Table 9: Reduction in Mean Over Placebo in Weekly Frequency Of Partial Onset Seizures in Study 5

|   | Placebo (N=97) | Immediate-Release Levetiracetam Tablets (N=101) |
|---|----------------|---|
| Percent reduction in partial seizure frequency over placebo | –              | 26.8%*  |

\*statistically significant versus placebo

The percentage of patients (y-axis) who achieved ≥ 50% reduction in weekly partial onset seizure frequency over the entire randomized treatment period (titration + evaluation period) within the two treatment groups (x-axis) in Study 5 is presented in Figure 4.

Figure 4: Responder Rate (≥ 50% Reduction From Baseline) in Study 5



\* statistically significant versus placebo

## 16 HOW SUPPLIED/STORAGE AND HANDLING

### 16.1 How Supplied

Levetiracetam extended-release tablets, USP 500 mg are white, modified oval-shaped, convex, film-coated, debossed "4186" on one side and debossed "V" on the reverse side. They are available as follows:

Bottles of 60 NDC 0603-4186-20

Levetiracetam extended-release tablets, USP, 750 mg are white, modified oval-shaped, convex, film-coated, debossed "4187" on one side and debossed "V" on the reverse side. They are available as follows:

Bottles of 60 NDC 0603-4187-20

### 16.2 Storage

Store at 20° to 25°C (68° to 77°F) (see USP Controlled Room Temperature).

## 17 PATIENT COUNSELING INFORMATION

Advise the patient to read the FDA-approved Patient Labeling (Medication Guide).

### Suicidal Behavior and Ideation

Counsel patients, their caregivers, and/or families that antiepileptic drugs (AEDs), including levetiracetam extended-release tablets, may increase the risk of suicidal thoughts and behavior and advise patients to be alert for the emergence or worsening of symptoms of depression; unusual changes in mood or behavior; or suicidal thoughts, behavior, or thoughts about self-harm. Advise patients, their caregivers, and/or families to immediately report behaviors of concern to a healthcare provider.

### Psychiatric Reactions and Changes in Behavior

Advise patients that levetiracetam extended-release tablets may cause changes in behavior (e.g. irritability and aggression). In addition, patients should be advised that they may experience changes in behavior that have been seen with other formulations of levetiracetam, which include agitation, anger, anxiety, apathy, depression, hostility, and psychotic symptoms.

### Effects on Driving or Operating Machinery

**MEDICATION GUIDE**  
**LEVETIRACETAM EXTENDED-RELEASE TABLETS, USP**  
 (lee" va tye ra' se tam)  
 500 mg and 750 mg

**Rx only**

Read this Medication Guide before you start taking levetiracetam extended-release tablets and each time you get a refill. There may be new information. This information does not take the place of talking to your healthcare provider about your medical condition or treatment.

**What is the most important information I should know about levetiracetam extended-release tablets?**

**Like other antiepileptic drugs, levetiracetam extended-release tablets may cause suicidal thoughts or actions in a very small number of people, about 1 in 500 people taking it.**

**Call a healthcare provider right away if you have any of these symptoms, especially if they are new, worse, or worry you:**

- thoughts about suicide or dying
- attempts to commit suicide
- new or worse depression
- new or worse anxiety
- feeling agitated or restless
- panic attacks
- trouble sleeping (insomnia)
- new or worse irritability
- acting aggressive, being angry, or violent
- acting on dangerous impulses
- an extreme increase in activity and talking (mania)
- other unusual changes in behavior or mood

**Do not stop levetiracetam extended-release tablets without first talking to a healthcare provider.**

- Stopping levetiracetam extended-release tablets suddenly can cause serious problems. Stopping a seizure medicine suddenly can cause seizures that will not stop (status epilepticus).
- Suicidal thoughts or actions can be caused by things other than medicines. If you have suicidal thoughts or actions, your healthcare provider may check for other causes.

**How can I watch for early symptoms of suicidal thoughts and actions?**

- Pay attention to any changes, especially sudden changes, in mood, behaviors, thoughts, or feelings.
- Keep all follow-up visits with your healthcare provider as scheduled.
- Call your healthcare provider between visits as needed, especially if you are worried about symptoms.

**What are levetiracetam extended-release tablets?**

Levetiracetam extended-release tablets are a prescription medicine taken by mouth that is used with other medicines to treat partial onset seizures in people 12 years of age and older with epilepsy.

It is not known if levetiracetam extended-release tablets are safe or effective in people under 12 years of age.

Before taking your medicine, make sure you have received the correct medicine. Compare the name above with the name on your bottle and the appearance of your medicine with the description of levetiracetam extended-release tablets provided below. Tell your pharmacist immediately if you think you have been given the wrong medicine.

500 mg levetiracetam extended-release tablets are white, modified oval-shaped, convex, film-coated, debossed "4186" on one side and debossed "V" on the reverse side.

750 mg levetiracetam extended-release tablets are white, modified oval-shaped, convex, film-coated, debossed "4187" on one side and debossed "V" on the reverse side.

**What should I tell my healthcare provider before starting levetiracetam extended-release tablets?**

Before taking levetiracetam extended-release tablets, tell your healthcare provider about all of your medical conditions, including if you:

- have or have had depression, mood problems or suicidal thoughts or behavior
- have kidney problems
- are pregnant or planning to become pregnant. It is not known if levetiracetam extended-release tablets will harm your unborn baby. You and your healthcare provider will have to decide if you should take levetiracetam extended-release tablets while you are pregnant. If you become pregnant while taking levetiracetam extended-release tablets, talk to your healthcare provider about registering with the North American Antiepileptic Drug Pregnancy Registry. You can enroll in this registry by calling 1-888-233-2334. The purpose of this registry is to collect information about the safety of levetiracetam extended-release tablets and other antiepileptic medicine during pregnancy.
- are breast feeding. Levetiracetam can pass into your milk and may harm your baby. You and your healthcare provider should discuss whether you should take levetiracetam extended-release tablets or breast feed; you should not do both.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Do not start a new medicine without first talking with your healthcare provider.

Know the medicines you take. Keep a list of them to show your healthcare provider and pharmacist each time you get a new medicine.

**How should I take levetiracetam extended-release tablets?**

Take levetiracetam extended-release tablets exactly as prescribed.

- Your healthcare provider will tell you how much levetiracetam extended-release tablets to take and when to take it. Levetiracetam extended-release tablets are usually taken once a day. Take levetiracetam extended-release tablets at the same time each day.
- Your healthcare provider may change your dose. Do not change your dose without talking to your healthcare provider.
- Take levetiracetam extended-release tablets with or without food.
- Swallow the tablets whole. Do not chew, break, or crush tablets.
- The inactive part of levetiracetam extended-release tablets may not dissolve after all the medicine has been released in your body. You may sometimes notice something in your bowel movement that looks like swollen pieces of the original tablet. This is normal.
- If you miss a dose of levetiracetam extended-release tablets, take it as soon as you remember. If it is almost time for your next dose, just skip the missed dose. Take the next dose at your regular time. **Do not take two doses at the same time.**
- If you take too much levetiracetam extended-release tablets, call your local Poison Control Center or go to the nearest emergency room right away.

**What should I avoid while taking levetiracetam extended-release tablets?**

Do not drive, operate machinery or do other dangerous activities until you know how levetiracetam extended-release tablets affect you. Levetiracetam extended-release tablets may make you dizzy or sleepy.

**What are the possible side effects of levetiracetam extended-release tablets?**

- See "What is the most important information I should know about levetiracetam extended-release tablets?"

Levetiracetam extended-release tablets can cause serious side effects.

Call your healthcare provider right away if you have any of these symptoms:

- mood and behavior changes such as aggression, agitation, anger, anxiety, apathy, mood swings, depression, hostility, and irritability. A few people may get psychotic symptoms such as hallucinations (seeing or hearing things that are really not there), delusions (false or strange thoughts or beliefs) and unusual behavior.
- extreme sleepiness, tiredness, and weakness
- problems with muscle coordination (problems walking and moving)
- a skin rash. Serious skin rashes can happen after you start taking levetiracetam extended-release tablets. There is no way to tell if a mild rash will become a serious reaction.

Common side effects seen in people who take levetiracetam extended-release tablets and other formulations of levetiracetam include:

- sleepiness
- weakness
- infection
- dizziness

These side effects can happen at any time but happen more often within the first 4 weeks of treatment except for infection.

Tell your healthcare provider if you have any side effect that bothers you or that does not go away.

These are not all the possible side effects of levetiracetam extended-release tablets. For more information, ask your healthcare provider or pharmacist.

**Call your doctor for medical advice about side effects. You may also report side effects to FDA at 1-800-FDA-1088.**

**How should I store levetiracetam extended-release tablets?**

- Store levetiracetam extended-release tablets at room temperature, 68°F to 77°F (20°C to 25°C) away from heat and light.
- **Keep levetiracetam extended-release tablets and all medicines out of the reach of children.**

**General information about the safe and effective use of levetiracetam extended-release tablets.**

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use levetiracetam extended-release tablets for a condition for which it was not prescribed. Do not give levetiracetam extended-release tablets to other people, even if they have the same symptoms that you have. It may harm them.

This Medication Guide summarizes the most important information about levetiracetam extended-release tablets. If you would like more information, talk with your healthcare provider. You can ask your pharmacist or healthcare provider for information about levetiracetam extended-release tablets that is written for health professionals.

**What are the ingredients of levetiracetam extended-release tablets?**

**Levetiracetam extended-release tablet** active ingredient: levetiracetam

Inactive ingredients: colloidal silicon dioxide, hydroxypropyl cellulose, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, pregelatinized starch (maize), stearic acid, titanium dioxide, and triacetin.

Levetiracetam extended-release tablets do not contain gluten.

This Medication Guide has been approved by the U.S. Food and Drug Administration.

Manufactured for:  
**QUALITEST PHARMACEUTICALS**  
 Huntsville, AL 35811

8183227  
 Revised: 5/2016  
 R4

Part 2 of 2  
 Rx only  
 500 mg and 750 mg  
 Tablets  
 Extended-Release  
 Levetiracetam



8183226 R7  
 Part 2 of 2

*The Challenge Printing Company*

|   |                      |                                   |                           |                          |             |
|---|----------------------|-----------------------------------|---------------------------|--------------------------|-------------|
|  | Customer: Par        | Flat Size: 12.25" x 11.25"        | Other:                    | <b>CUSTOMER APPROVAL</b> | <b>DATE</b> |
|   | Proof ID#: 223863-1  | Fold Size: 1.25" x 1.25"          | <b>CHALLENGE APPROVAL</b> | <b>INITIALS</b>          | <b>DATE</b> |
|   | Proof Date: 05/17/16 | Perforation: N/A                  | ARTWORK SET BY:           |                          |             |
|   | Colors: Black        | Outsert Orientation: Single Sided | QA APPROVAL BY:           |                          |             |

NOTE(S): 1. This electronic proof, for your written approval, is provided for purposes of indicating color break and image placement only, and not as a representation of color fidelity.

2. Barcode(s) provided on this proof is (are) for purposes of indicating encodation only and not to be used for grading according to ANSI standards, as applicable.