HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use OLANZAPINE and FLUOXETINE CAPSULES, safely and effectively. See full prescribing information for OLANZAPINE and FLUOXETINE CAPSULES.

OLANZAPINE and FLUOXETINE CAPSULES, for oral use

Initial U.S. Approval: 2003

WARNING: SUICIDAL THOUGHTS AND BEHAVIORS; AND INCREASED MORTALITY IN ELDERLY PATIENTS WITH DEMENTIA-RELATED PSYCHOSIS

See full prescribing information for complete boxed warning.

- **Increased risk of suicidal thinking and behavior in children, adolescents, and young adults taking antidepressants. Olanzapine and Fluoxetine Capsules are not approved for use in children less than 10 years of age (5.1, 8.4, 17.2).**
- **Monitor for worsening and emergence of suicidal thoughts and behaviors (5.1, 17.2).**
- **Elderly patients with dementia-related psychosis treated with antipsychotic drugs are at an increased risk of death. Olanzapine and Fluoxetine Capsules are not approved for the treatment of patients with dementia-related psychosis (5.2, 5.21, 17.3).**

---------- INDICATIONS AND USAGE ----------

Olanzapine and Fluoxetine Capsules combines olanzapine, an atypical antipsychotic and fluoxetine, a selective serotonin reuptake inhibitor, indicated for treatment of:

- Acute Depressive Episodes Associated with Bipolar I Disorder (1)
- Treatment Resistant Depression (1)

---------- DOSAGE AND ADMINISTRATION ----------

- Adult Starting Dose: 6 mg olanzapine with 25 mg fluoxetine (6 mg/25 mg, once daily in the evening (**2.1, 2.2**)
- Adult Maximum Dose: 12 mg/50 mg once daily (**2.1, 2.2**)
- Pediatric Bipolar Depression Starting Dose: 3 mg/25 mg once daily (for ages 10 to 17 years) (**2.1**).
- Pediatric Bipolar Depression Maximum Dose: 12 mg/50 mg (**2.1**).
- Starting dose in patients predisposed to hypotensive reactions, hepatic impairment, or with potential for slowed metabolism: 3 mg/25 mg to 6 mg /25 mg. Escalate dose cautiously (**2.3**)

---------- DOSAGE FORMS AND STRENGTHS ----------

- Capsules: 3 mg/25 mg, 6 mg/25 mg, 6 mg/50 mg, 12 mg/25 mg, and 12 mg/50 mg (mg olanzapine/mg equivalent fluoxetine) (**3**)

---------- CONTRAINDICATIONS ----------

- *Monoamine Oxidase Inhibitors (MAOI)*: Because of the risk of serotonin syndrome, do not use MAOIs intended to treat psychiatric disorders with Olanzapine and Fluoxetine Capsules or within 5 weeks of stopping treatment with Olanzapine and Fluoxetine Capsules. Do not use Olanzapine and Fluoxetine Capsules within 14 days of stopping an MAOI intended to treat psychiatric disorders. In addition, do not start Olanzapine and Fluoxetine Capsules in a patient who is being treated with linezolid or intravenous methylene blue. (**4.1**)
- *Pimozide*: Do not use. Risk of QT interval prolongation (**4.2, 5.20, 7.7, 7.8**)
- *Thioridazine*: Do not use. Risk of QT interval prolongation. Do not use thioridazine within 5 weeks of discontinuing Olanzapine and Fluoxetine Capsules (**4.2, 5.20, 7.7, 7.8**)

---------- WARNINGS AND PRECAUTIONS ----------

- *Neuroleptic Malignant Syndrome*: Manage with immediate discontinuation and close monitoring (**5.3**)
- *Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS)*: Discontinue if DRESS is suspected (**5.4**)
- *Metabolic Changes*: Atypical antipsychotic drugs have been associated with metabolic changes including hyperglycemia, dyslipidemia, and weight gain (**5.5**)
  - *Hyperglycemia and Diabetes Mellitus*: In some cases extreme and associated with ketoacidosis or hyperosmolar coma or death. Monitor for symptoms of hyperglycemia. Perform fasting blood glucose testing before beginning, and periodically during treatment. (**5.5**)
  - *Dyslipidemia*: Appropriate clinical monitoring is recommended, including fasting blood lipid testing before beginning, and periodically during, treatment (**5.5**)
  - *Weight Gain*: Consider potential consequences of weight gain. Monitor weight regularly (**5.5**)
- *Serotonin Syndrome*: Serotonin syndrome has been reported with SSRIs and SNRIs, including Olanzapine and Fluoxetine Capsules, both when taken alone, but especially when coadministered with other serotonergic agents (including triptans, tricyclic antidepressants, fentanyl, lithium, tramadol, tryptophan, buspirone, amphetamines and St. John's Wort), and with drugs that impair metabolism of serotonin (in particular, MAOIs, both those intended to treat psychiatric disorders and others, such as linezolid and intravenous methylene blue). Serotonin syndrome symptoms may include mental status changes (e.g., agitation, hallucinations, delirium, and coma), autonomic instability (e.g., tachycardia, labile blood pressure, dizziness, diaphoresis, flushing, hyperthermia), neuromuscular symptoms (e.g., tremor, rigidity, myoclonus, hyperreflexia, incoordination), seizures, and/or gastrointestinal symptoms (e.g., nausea, vomiting, diarrhea). Patients should be monitored for the emergence of serotonin syndrome.

  The concomitant use of olanzapine and fluoxetine with MAOIs intended to treat psychiatric disorders is contraindicated. Olanzapine and Fluoxetine should also not be started in a patient who is being treated with MAOIs such as linezolid or intravenous methylene blue. All reports with methylene blue that provided information on the route of administration involved intravenous administration in the dose range of 1mg/kg to 8mg/kg. No reports involved the administration of methylene blue by other routes (such as oral tablets or local tissue injection) or at lower doses. There may be circumstances when it is necessary to initiate treatment with an MAOI such as linezolid or intravenous methylene blue in a patient taking olanzapine and fluoxetine. Olanzapine and fluoxetine should be discontinued before initiating treatment with the MAOI *[see Dosage and Administration (2.4, 2.5) and Contraindications (4.1)]*.

  If concomitant use of olanzapine and fluoxetine with other serotonergic drugs including triptans, tricyclic antidepressants, fentanyl, lithium, tramadol, buspirone, tryptophan, amphetamines, and St. John's Wort is clinically warranted, patients should be made aware of a potential increased risk for serotonin syndrome, particularly during treatment initiation and dose increases (**5.6**).
- *Angle-Closure Glaucoma*: Angle-closure glaucoma has occurred in patients with untreated anatomically narrow angles treated with antidepressants (**5.7**)
- *Allergic Reactions and Rash*: Discontinue upon appearance of rash or allergic phenomena (**5.8**)
- *Activation of Mania/Hypomania*: Screen for Bipolar Disorder and monitor for activation of mania/hypomania (**5.9**)
- *Fluoxetine Withdrawal Syndrome* if clinically appropriate (**5.10**)
- *Orthostatic Hypotension*: Can be associated with bradycardia and syncope. Risk is increased during initial dose titration. Use caution in patients with cardiovascular disease or cerebrovascular disease, and those conditions that could affect hemodynamic responses (**5.11**)
- *Leukopenia, Neutropenia, and Agranulocytosis*: Has been reported with antipsychotics, including Olanzapine and Fluoxetine. Patients with a history of a clinically significant low white blood cell count (WBC) or drug induced leukopenia/neutropenia should have their complete blood count (CBC) monitored frequently during the first few months of therapy. Consider discontinuing Olanzapine and Fluoxetine Capsules at the first sign of a clinically significant decline in WBC in the absence of other causative factors (**5.12**)
- *Seizures*: Use cautiously in patients with a history of seizures or with conditions that lower the seizure threshold (**5.15**)
- *Abnormal Bleeding*: SSRIs increase the risk of bleeding. Use with NSAIDs, aspirin, warfarin, or other drugs that affect coagulation may potentiate the risk of gastrointestinal or other bleeding (**5.16**)
- *Hyponatremia*: Can occur in association with syndrome of inappropriate antidiuretic hormone (SIADH). Consider discontinuing Olanzapine and Fluoxetine capsules if symptomatic hyponatremia occurs (SIADH) (**5.17**)
- *Potential for Cognitive and Motor Impairment*: Has potential to impair judgment, thinking, and motor skills. Caution patients about operating machinery (**5.18**)
- *QT Prolongation*: QT prolongation and ventricular arrhythmia including Torsade de Pointes have been reported with fluoxetine. Use with caution in conditions that predispose to arrhythmias or increased fluoxetine exposure. Use cautiously in patients with risk factors for QT prolongation (**4.2, 5.20**)
- *Hyperprolactinemia*: May elevate prolactin levels (**5.22**)
- *Long Elimination Half-Life of Fluoxetine*: Changes in dose will not be fully reflected in plasma for several weeks (**5.24**)

---------- ADVERSE REACTIONS ----------

Most common adverse reactions (≥5% and at least twice that for placebo) in adults: sedation, weight increased, appetite increased, dry mouth, fatigue, edema, tremor, disturbance in attention, blurred vision. Children and adolescents: sedation, weight increased, appetite increased, tremor, triglyceride increased, hepatic enzymes increased (**6.1**)

To report SUSPECTED ADVERSE REACTIONS, contact Par Pharmaceutical at 1-800-828-9393 or FDA at 1-800- FDA-1088 or www.fda.gov/medwatch.

---------- DRUG INTERACTIONS ----------

- *Monoamine Oxidase Inhibitor (MAOI)*: (**2.4, 2.5, 4.1, 5.6, 7.1**)
- *Drugs Metabolized by CYP2D6*: Fluoxetine is a potent inhibitor of CYP2D6 enzyme pathway (**7.7**)
- *Tricyclic Antidepressants (TCAs)*: Monitor TCA levels during coadministration with olanzapine and fluoxetine or when olanzapine and fluoxetine capsules have been recently discontinued (**5.6, 7.7**)
- *CNS Acting Drugs*: Caution is advised if the concomitant administration of olanzapine and fluoxetine with other CNS-active drugs is required (**7.2**)
- *Antihypertensive Agent*: Enhanced antihypertensive effect (**7.7**)
- *Levodopa and Dopamine Agonists*: May antagonize levodopa/dopamine agonists (**7.7**)
- *Benzodiazepines*: May potentiate orthostatic hypotension and sedation (**7.6, 7.7**)
- *Clozapine*: May elevate clozapine levels (**7.7**)
- *Haloperidol*: Elevated haloperidol levels have been observed (**7.7**)
- *Carbamazepine*: Potential for elevated carbamazepine levels and clinical anticonvulsant toxicity (**7.7**)
- *Phenytoin*: Potential for elevated phenytoin levels and clinical anticonvulsant toxicity (**7.7**)
- *Alcohol*: May potentiate sedation and orthostatic hypotension (**7.7**)
- *Serotonergic Drugs*: (**2.4, 2.5, 4.1, 5.6, 7.3**)
- *Fluvoxamine*: May increase olanzapine levels; a lower dose of the olanzapine component of olanzapine and fluoxetine capsules should be considered (**7.8**)
- *Drugs that Interfere with Hemostasis (e.g., NSAIDs, Aspirin, Warfarin, etc.)*: May potentiate the risk of bleeding (**7.4**)
- *Drugs Tightly Bound to Plasma Proteins*: Fluoxetine may cause shift in plasma concentrations (**7.7**)
- *Drugs that Prolong the QT Interval*: Do not use olanzapine and fluoxetine capsules in combination with thioridazine or pimozide. Use olanzapine and fluoxetine capsules with caution in combination with other drugs that prolong the QT interval (**4.2, 5.20, 7.7, 7.8**)

---------- USE IN SPECIFIC POPULATIONS ----------

- *Pregnancy*: Olanzapine and Fluoxetine capsules should be given during pregnancy only if the potential benefit justifies the potential risk to the fetus (**8.1**)
- *Nursing Mothers*: Breast feeding is not recommended (**8.3**)
- *Pediatric Use*: Safety and efficacy of olanzapine and fluoxetine capsules for the treatment of bipolar I depression in patients under 10 years of age have not been established. Safety and efficacy of olanzapine and fluoxetine capsules for treatment resistant depression in patients under 18 years of age have not been established (**8.4**).
- *Hepatic Impairment*: Use a lower or less frequent dose in patients with cirrhosis (**8.6**)

See 17 for PATIENT COUNSELING INFORMATION and the FDA-approved Medication Guide

Revised: 05/2019

FULL PRESCRIBING INFORMATION: CONTENTS *

WARNINGS: SUICIDAL THOUGHTS AND BEHAVIORS; AND INCREASED MORTALITY IN ELDERLY PATIENTS WITH DEMENTIA-RELATED PSYCHOSIS

1 INDICATIONS AND USAGE
2 DOSAGE AND ADMINISTRATION
- 2.1 Depressive Episodes Associated with Bipolar I Disorder
- 2.2 Treatment Resistant Depression
- 2.3 Specific Populations
- 2.4 Switching a Patient To or From a Monoamine Oxidase Inhibitor (MAOI) Intended to Treat Psychiatric Disorders
- 2.5 Use of Olanzapine and Fluoxetine Capsules with other MAOIs such as Linezolid or Methylene Blue

2.6 Discontinuation of Treatment with Olanzapine and Fluoxetine Capsules

3 DOSAGE FORMS AND STRENGTHS
4 CONTRAINDICATIONS
- 4.1 Monoamine Oxidase Inhibitors (MAOIs)
- 4.2 Other Contraindications

5 WARNINGS AND PRECAUTIONS
- 5.1 Suicidal Thoughts and Behaviors in Children, Adolescents, and Young Adults
- 5.2 Increased Mortality in Elderly Patients with Dementia-Related Psychosis
- 5.3 Neuroleptic Malignant Syndrome (NMS)
- 5.4 Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS)
- 5.5 Metabolic Changes
- 5.6 Serotonin Syndrome
- 5.7 Angle-Closure Glaucoma
- 5.8 Allergic Reactions and Rash
- 5.9 Activation of Mania/Hypomania
- 5.10 Tardive Dyskinesia
- 5.11 Orthostatic Hypotension
- 5.12 Falls
- 5.13 Leukopenia, Neutropenia, and Agranulocytosis
- 5.14 Hyponatremia
- 5.15 Seizures
- 5.16 Abnormal Bleeding
- 5.17 Hyperprolactinemia
- 5.18 Potential for Cognitive and Motor Impairment
- 5.19 Body Temperature Dysregulation
- 5.20 QT Prolongation
- 5.21 Use in Patients with Concomitant Illness
- 5.22 Hyperprolactinemia
- 5.23 Concomitant Use of Olanzapine and Fluoxetine Products
- 5.24 Long Elimination Half-Life of Fluoxetine
- 5.25 Discontinuation Adverse Reactions

6 ADVERSE REACTIONS
- 6.1 Clinical Trials Experience
- 6.2 Vital Sign and Laboratory Studies
- 6.3 Postmarketing Experience

7 DRUG INTERACTIONS
- 7.1 Monoamine Oxidase Inhibitors (MAOIs)
- 7.2 CNS Acting Drugs
- 7.3 Serotonergic Drugs
- 7.4 Drugs that Interfere with Hemostasis (e.g., NSAIDs, Aspirin, Warfarin)
- 7.5 Electroconvulsive Therapy (ECT)
- 7.6 Potential for Other Drugs to Affect Olanzapine and Fluoxetine
- 7.7 Potential for Olanzapine and Fluoxetine to Affect Other Drugs
- 7.8 Drugs that Prolong the QT Interval

8 USE IN SPECIFIC POPULATIONS
- 8.1 Pregnancy
- 8.2 Labor and Delivery
- 8.3 Nursing Mothers
- 8.4 Pediatric Use
- 8.5 Geriatric Use
- 8.6 Hepatic Impairment

9 DRUG ABUSE AND DEPENDENCE
- 9.3 Dependence

10 OVERDOSAGE
- 10.1 Management of Overdose

11 DESCRIPTION
12 CLINICAL PHARMACOLOGY
- 12.1 Mechanism of Action
- 12.2 Pharmacodynamics
- 12.3 Pharmacokinetics
- 12.4 Specific Populations

13 NONCLINICAL TOXICOLOGY
- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

14 CLINICAL STUDIES
- 14.1 Depressive Episodes Associated with Bipolar I Disorder
- 14.2 Treatment Resistant Depression

16 HOW SUPPLIED/STORAGE AND HANDLING
- 16.1 How Supplied
- 16.2 Storage and Handling

17 PATIENT COUNSELING INFORMATION
- 17.1 Information with Medication Guide
- 17.2 Suicidal Thoughts and Behaviors in Children, Adolescents, and Young Adults
- 17.3 Elderly Patients with Dementia-Related Psychosis: Increased Mortality and Cerebrovascular Adverse Events (CVAE), Including Stroke
- 17.4 Neuroleptic Malignant Syndrome (NMS)
- 17.5 Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS)
- 17.6 Hyperglycemia and Diabetes Mellitus
- 17.7 Dyslipidemia
- 17.8 Weight Gain
- 17.9 Serotonin Syndrome
- 17.10 Angle-Closure Glaucoma
- 17.11 Allergic Reactions and Rash
- 17.12 Orthostatic Hypotension
- 17.13 Abnormal Bleeding
- 17.14 Hyponatremia
- 17.15 Potential for Cognitive and Motor Impairment
- 17.16 Body Temperature Dysregulation
- 17.17 Concomitant Medication
- 17.18 Discontinuation of Treatment with Olanzapine and Fluoxetine Capsules
- 17.19 Alcohol
- 17.20 Use in Specific Populations
- 17.21 QT Prolongation

* Sections or subsections omitted from the full prescribing information are not listed

FULL PRESCRIBING INFORMATION

WARNINGS: SUICIDAL THOUGHTS AND BEHAVIORS; AND INCREASED MORTALITY IN ELDERLY PATIENTS WITH DEMENTIA-RELATED PSYCHOSIS

Suicidal Thoughts and Behaviors – Antidepressants increased the risk of suicidal thoughts and behavior in children, adolescents, and young adults in short-term studies. These studies did not show an increase in the risk of suicidal thoughts and behavior with antidepressant use in patients over age 24; there was a reduction in risk with antidepressant use in patients aged 65 and older.

In patients of all ages who are started on antidepressant therapy, monitor closely for worsening and emergence of suicidal thoughts and behaviors. Advise families and caregivers of the need for close observation and communication with the prescriber. Olanzapine and Fluoxetine Capsules are not approved for use in children less than 10 years of age *[see Warnings and Precautions (5.1), Use in Specific Populations (8.4), and Patient Counseling Information (17.2)]*.

Increased Mortality in Elderly Patients with Dementia-Related Psychosis — Elderly patients with dementia-related psychosis treated with antipsychotic drugs are at an increased risk of death. Olanzapine and Fluoxetine Capsules are not approved for the treatment of patients with dementia-related psychosis *[see Warnings and Precautions (5.2)]*.

1 INDICATIONS AND USAGE

Olanzapine and Fluoxetine is indicated for the treatment of:

- Acute depressive episodes in Bipolar I Disorder *[see Clinical Studies (14.1)]*.
- Treatment resistant depression (Major Depressive Disorder in patient who do not respond to 2 separate trials of different antidepressants of adequate dose and duration in the current episode) *[see Clinical Studies (14.2)]*.

2 DOSAGE AND ADMINISTRATION

2.1 Depressive Episodes Associated with Bipolar I Disorder

Adults – Administer Olanzapine and Fluoxetine Capsules once daily in the evening, generally beginning with the 6 mg/25 mg (mg olanzapine/mg equivalent fluoxetine) capsule. While food has no appreciable effect on the absorption of olanzapine and fluoxetine given individually, the effect of food on the absorption of olanzapine and fluoxetine has not been studied. Make dosage adjustments, if indicated, according to efficacy and tolerability. Antidepressant efficacy was demonstrated with olanzapine and fluoxetine in a dose range of olanzapine 6 mg to 12 mg and fluoxetine 25 mg to 50 mg *[see Clinical Studies (14.1)]*. The safety of doses above 18 mg of olanzapine and 75 mg of fluoxetine has not been evaluated in adult clinical studies. Periodically reexamine the need for continued pharmacotherapy.

Children and Adolescents (10 to 17 years of age) – Administer Olanzapine and Fluoxetine Capsules once daily in the evening, generally beginning with the 3 mg/25 mg capsule, without regard to meals, with a recommended target dose within the approved dosing range (6/26, 6/50, 12/25, 12/50 mg) *[see Clinical Studies (14.1)]*. The safety of doses above 12 mg of olanzapine and 50 mg of fluoxetine has not been evaluated in pediatric clinical studies. Periodically reexamine the need for continued pharmacotherapy.

2.2 Treatment Resistant Depression

Administer Olanzapine and Fluoxetine Capsules once daily in the evening, generally beginning with the 6 mg/25 mg capsule. While food has no appreciable effect on the absorption of olanzapine and fluoxetine given individually, the effect of food on the absorption of Olanzapine and Fluoxetine Capsules has not been studied. Adjust dosage, if indicated, according to efficacy and tolerability.

Antidepressant efficacy was demonstrated with Olanzapine and Fluoxetine Capsules in a dose range of olanzapine 6 mg to 18 mg and fluoxetine 25 mg to 50 mg *[see Clinical Studies (14.2)]*. The safety of doses above 18 mg/75 mg has not been evaluated in clinical studies. Periodically reexamine the need for continued pharmacotherapy.

2.3 Specific Populations

Start Olanzapine and Fluoxetine Capsules 3 mg/25 mg or 6 mg/25 mg in patients with a predisposition to hypotensive reactions, patients with hepatic impairment, or patients who exhibit a combination of factors that may slow the metabolism of olanzapine and fluoxetine (female gender, geriatric age, nonsmoking status) or those patients who may be pharmacodynamically sensitive to olanzapine. Titrate slowly and adjust dosage as needed in patients who exhibit a combination of factors that may slow metabolism. Olanzapine and Fluoxetine Capsules have not been systematically studied in patients >65 years of age or in patients <10 years of age *[see Warnings and Precautions (5.21), Use in Specific Populations (8.5), and Clinical Pharmacology (12.3, 12.4)]*.

Treatment of Pregnant Women– When treating pregnant women with fluoxetine, a component of olanzapine and fluoxetine, the physician should carefully consider the potential risks and potential benefits of treatment. Neonates exposed to SSRIs or SNRIs late in the third trimester have developed complications requiring prolonged hospitalization, respiratory support, and tube feeding *[see Use in Specific Populations (8.1)]*.

2.4 Switching a Patient To or From a Monoamine Oxidase Inhibitor (MAOI) Intended to Treat Psychiatric Disorders

At least 14 days should elapse between discontinuation of an MAOI intended to treat psychiatric disorders and initiation of therapy with olanzapine and fluoxetine. Conversely, at least 5 weeks should be allowed after stopping olanzapine and fluoxetine before starting an MAOI intended to treat psychiatric disorders *[see Contraindications (4.1)]*.

2.5 Use of Olanzapine and Fluoxetine Capsules with Other MAOIs such as Linezolid or Methylene Blue

Do not start olanzapine and fluoxetine in a patient who is being treated with linezolid or intravenous methylene blue because there is an increased risk of serotonin syndrome. In a patient who requires more urgent treatment of a psychiatric condition, other interventions, including hospitalization, should be considered *[see Contraindications (4.1)]*.

In some cases, a patient already receiving olanzapine and fluoxetine therapy may require urgent treatment with linezolid or intravenous methylene blue. If acceptable alternatives to linezolid or intravenous methylene blue treatment are not available and the potential benefits of linezolid or intravenous methylene blue are judged to outweigh the risks of serotonin syndrome in a particular patient, olanzapine and fluoxetine should be stopped promptly, and linezolid or intravenous methylene blue can be administered. The patient should be monitored for symptoms of serotonin syndrome for five weeks or until 24 hours after the last dose of linezolid or intravenous methylene blue, whichever comes first. Therapy with olanzapine and fluoxetine may be resumed 24 hours after the last dose of linezolid or intravenous methylene blue *[see Warnings and Precautions (5.6)]*.

The risk of administering methylene blue by non-intravenous routes (such as oral tablets or by local injection) or in intravenous doses much lower than 1 mg/kg with olanzapine and fluoxetine is unclear. The clinician should, nevertheless, be aware of the possibility of emergent symptoms of serotonin syndrome with such use *[see Warnings and Precautions (5.6)]*.

2.6 Discontinuation of Treatment with Olanzapine and Fluoxetine Capsules

Symptoms associated with discontinuation of fluoxetine, a component of olanzapine and fluoxetine, SNRIs, and SSRIs, have been reported *[see Warnings and Precautions (5.25)]*.

3 DOSAGE FORMS AND STRENGTHS

Capsules (mg olanzapine/mg equivalent fluoxetine):

- 3 mg/25 mg
- 6 mg/25 mg
- 6 mg/50 mg
- 12 mg/25 mg
- 12 mg/50 mg

4 CONTRAINDICATIONS

4.1 Monoamine Oxidase Inhibitors (MAOIs)

The use of MAOIs intended to treat psychiatric disorders with olanzapine and fluoxetine or within 5 weeks of stopping treatment with olanzapine and fluoxetine is contraindicated because of an increased risk of serotonin syndrome. The use of olanzapine and fluoxetine within 14 days of stopping an MAOI intended to treat psychiatric disorders is also contraindicated *[see Dosage and Administration (2.4) and Warnings and Precautions (5.6)]*.

Starting olanzapine and fluoxetine in a patient who is being treated with MAOIs such as linezolid or intravenous methylene blue is also contraindicated because of an increased risk of serotonin syndrome *[see Dosage and Administration (2.5) and Warnings and Precautions (5.6)]*.

4.2 Other Contraindications

- Pimozide — *[see Warnings and Precautions (5.20) and Drug Interactions (7.7, 7.8)]*
- Thioridazine — *[see Warnings and Precautions (5.20) and Drug Interactions (7.7, 7.8)]*

Pimozide and thioridazine prolong the QT interval. Olanzapine and fluoxetine are also known to prolong the levels of pimozide and thioridazine through inhibition of CYP2D6. Olanzapine and Fluoxetine can also prolong the QT interval.

5 WARNINGS AND PRECAUTIONS

5.1 Suicidal Thoughts and Behaviors in Children, Adolescents, and Young Adults

Patients with Major Depressive Disorder (MDD), both adult and pediatric, may experience worsening of their depression and/or the emergence of suicidal ideation and behavior (suicidality) or unusual changes in behavior, whether or not they are taking antidepressant medications, and this risk may persist until significant remission occurs. Suicide is a known risk of depression and certain other psychiatric disorders, and these disorders themselves are the strongest predictors of suicide. There has been a long-standing concern, however, that antidepressants may have a role in inducing worsening of depression and the emergence of suicidality in certain patients during the early phases of treatment. Pooled analyses of short-term placebo-controlled trials of antidepressant drugs (SSRIs and others) showed that these drugs increase the risk of suicidal thinking and behavior (suicidality) in children, adolescents, and young adults (ages 18 to 24) with Major Depressive Disorder (MDD) and other psychiatric disorders. Short-term studies did not show an increase in the risk of suicidality with antidepressants compared to placebo in adults beyond age 24; there was a reduction with antidepressants compared to placebo in adults aged 65 and older.

The pooled analyses of placebo-controlled trials in children and adolescents with MDD, Obsessive Compulsive Disorder (OCD), or other psychiatric disorders included a total of 24 short-term trials of 9 antidepressant drugs in over 4400 patients. The pooled analyses of placebo-controlled trials in adults with MDD or other psychiatric disorders included a total of 295 short-term trials (median duration of 2 months) of 11 antidepressant drugs in over 77,000 patients. There was considerable variation in risk of suicidality among drugs, but a tendency toward an increase in the younger patients for almost all drugs studied. There were differences in absolute risk of suicidality across the different indications, with the highest incidence in MDD. The risk differences (drug versus placebo), however, were relatively stable within age strata and across indications. These risk differences (drug-placebo difference in the number of cases of suicidality per 1000 patients treated) are provided in **Table 1**.

Table 1: Suicidality per 1000 Patients Treated

Age Range	Drug-Placebo Difference in Number of Cases of Suicidality per 1000 Patients Treated
<18	Increases compared to Placebo 14 additional cases
18 to 24	5 additional cases
25 to 64	Decreases Compared to Placebo 1 fewer case
≥65	6 fewer cases

No suicides occurred in any of the pediatric trials. There were no suicides in the adult trials, but the number was not sufficient to reach any conclusion about drug effect on suicide.

It is unknown whether the suicidality risk extends to longer-term use, i.e., beyond several months. However, there is substantial evidence from placebo-controlled maintenance trials in adults with depression that the use of antidepressants can delay the recurrence of depression.

All patients being treated with antidepressants for any indication should be monitored appropriately and observed closely for clinical worsening, suicidality, and unusual changes in behavior, especially during the initial few months of a course of drug therapy, or at times of dose changes, either increases or decreases.

The following symptoms, anxiety, agitation, panic attacks, insomnia, irritability, hostility, aggressiveness, impulsivity, akathisia (psychomotor restlessness), hypomania, and mania, have been reported in adult and pediatric patients being treated with antidepressants for Major Depressive Disorder as well as for other indications, both psychiatric and nonpsychiatric. Although a causal link between the emergence of such symptoms and either the worsening of depression and/or the emergence of suicidal impulses has not been established, there is concern that such symptoms may represent precursors to emerging suicidality.

Consideration should be given to changing the therapeutic regimen, including possibly discontinuing the medication, in patients whose depression is persistently worse, or who are experiencing emergent suicidality or symptoms that might be precursors to worsening depression or suicidality, especially if these symptoms are severe, abrupt in onset, or were not part of the patient's presenting symptoms.

If the decision has been made to discontinue treatment, medication should be tapered as rapidly as is feasible, but with recognition that abrupt discontinuation can be associated with certain symptoms *[see Warnings and Precautions (5.25)]*.

Families and caregivers of patients being treated with antidepressants for Major Depressive Disorder or other indications, both psychiatric and nonpsychiatric, should be alerted about the need to monitor patients for the emergence of agitation, irritability, unusual changes in behavior, and the other symptoms described above, as well as the emergence of suicidality, and to report such symptoms immediately to healthcare providers. Such monitoring should include daily observation by families and caregivers. Prescriptions for olanzapine and fluoxetine should be written for the smallest quantity of capsules consistent with good patient management, in order to reduce the risk of overdose.

It should be noted that olanzapine and fluoxetine capsules are not approved for use in treating any indications in patients less than 10 years of age *[see Use in Specific Populations (8.4)]*.

5.2 Increased Mortality in Elderly Patients with Dementia-Related Psychosis

Elderly patients with dementia-related psychosis treated with antipsychotic drugs are at an increased risk of death. Olanzapine and Fluoxetine capsules are not approved for the treatment of patients with dementia-related psychosis *[see Boxed Warning and Warnings and Precautions (5.2)]*.

Olanzapine and Fluoxetine Combination Capsules

In a olanzapine-controlled clinical trials of elderly patients with dementia-related psychosis, the incidence of death in olanzapine-treated patients was significantly greater than placebo-treated patients (3.5% vs 1.5%, respectively).

Meta-Analysis of Antipsychotic Use in Dementia-Related Psychosis — Elderly patients with dementia-related psychosis treated with antipsychotic drugs are at an increased risk of death. Analyses of seventeen placebo-controlled trials (modal duration of 10 weeks), largely in patients taking atypical antipsychotic drugs, revealed a risk of death in drug-treated patients of between 1.6 to 1.7 times the risk of death in placebo-treated patients. Over the course of a typical 10-week controlled trial, the rate of death in drug-treated patients was about 4.5%, compared to a rate of about 2.6% in the placebo group. Although the causes of death were varied, most of the deaths appeared to be either cardiovascular (e.g., heart failure, sudden death) or infectious (e.g., pneumonia) in nature. Observational studies suggest that, similar to atypical antipsychotic drugs, treatment with conventional antipsychotic drugs may increase mortality. The extent to which the findings of increased mortality in observational studies may be attributed to the antipsychotic drug as opposed to some characteristic(s) of the patients is not clear. Olanzapine and Fluoxetine Capsules are not approved for the treatment of patients with dementia-related psychosis *[see Warnings and Precautions (5.21) and Patient Counseling Information (17.3)]*.

Cerebrovascular Adverse Events (CVAE), Including Stroke - Cerebrovascular adverse events (e.g., stroke, transient ischemic attack), including fatalities, were reported in 3 trials of olanzapine in elderly patients with dementia-related psychosis. In placebo-controlled trials, there was a significantly higher incidence of cerebrovascular adverse events in patients treated with olanzapine compared to patients treated with placebo. Olanzapine and olanzapine and fluoxetine capsules are not approved for the treatment of patients with dementia-related psychosis *[see Boxed Warning and Patient Counseling Information (17.3)]*.

5.3 Neuroleptic Malignant Syndrome (NMS)

A potentially fatal symptom complex sometimes referred to as NMS has been reported in association with administration of antipsychotic drugs, including olanzapine. Rhabdomyolysis, myoglobinuria, acute renal failure, muscle rigidity, altered mental status, and evidence of autonomic instability (irregular pulse or blood pressure, tachycardia, diaphoresis, and cardiac dysrhythmia). Additional signs may include elevated creatinine phosphokinase, myoglobinuria (rhabdomyolysis), and acute renal failure.

The diagnostic evaluation of patients with this syndrome is complicated. In arriving at a diagnosis, it is important to exclude cases where the clinical presentation includes both serious medical illness (e.g., pneumonia, systemic infection, etc.) and untreated or inadequately treated extrapyramidal signs and symptoms (EPS). Other important considerations in the differential diagnosis include central anticholinergic toxicity, heat stroke, drug fever, and primary central nervous system pathology.

The management of NMS should include: 1) immediate discontinuation of antipsychotic drugs and other drugs not essential to concurrent therapy, 2) intensive symptomatic treatment and medical monitoring, and 3) treatment of any concomitant serious medical problems for which specific treatments are available. There is no general agreement about specific pharmacological treatment regimens for NMS.

If after recovering from NMS, a patient requires treatment with an antipsychotic, the patient should be carefully monitored, since recurrences of NMS have been reported *[see Warnings and Precautions (5.5) and Patient Counseling Information (17.4, 17.9)]*.

5.4 Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS)

Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS) has been reported with olanzapine exposure. DRESS may present with a cutaneous reaction (such as rash or exfoliative dermatitis), eosinophilia, fever, and/or lymphadenopathy, with systemic complications such as hepatitis, nephritis, hematologic abnormalities, and/or pericarditis. DRESS is sometimes fatal. Discontinue olanzapine and fluoxetine capsules if DRESS is suspected *[see Patient Counseling Information (17.5)]*.

5.5 Metabolic Changes

Atypical antipsychotic drugs have been associated with metabolic changes including hyperglycemia, dyslipidemia, and weight gain. While all of the atypical antipsychotic drugs cause some degree of these metabolic changes, olanzapine's specific metabolic profile is presented below.

Hyperglycemia and Diabetes Mellitus

Adults - Physicians should consider the risks and benefits when prescribing olanzapine and fluoxetine capsule to patients with an established diagnosis of diabetes mellitus, or having borderline increased blood glucose level (fasting 100 to 126 mg/dL, or nonfasting 140 to 200 mg/dL). Fasting glucose and fluoxetine capsule levels should be monitored regularly for worsening of glucose control. Patients starting treatment with olanzapine and fluoxetine should undergo fasting blood glucose testing at the beginning of treatment and periodically during treatment. Any patient treated with atypical antipsychotics should be monitored for symptoms of hyperglycemia including polydipsia, polyuria, polyphagia, and weakness. Patients who develop symptoms of hyperglycemia during treatment with atypical antipsychotics should undergo fasting blood glucose testing. In some cases, hyperglycemia has resolved when the atypical antipsychotic was discontinued; however, some patients required continuation of anti-diabetic treatment despite discontinuation of the suspect drug *[see Patient Counseling Information (17.6)]*.

Hyperglycemia, in some cases extreme and associated with ketoacidosis or hyperosmolar coma or death, has been reported in patients treated with atypical antipsychotics, including olanzapine, as well as olanzapine taken concomitantly with fluoxetine. Assessment of the relationship between atypical antipsychotic use and glucose abnormalities is complicated by the possibility of an increased background risk of diabetes mellitus in patients with schizophrenia and the increasing incidence of diabetes mellitus in the general population. Epidemiological studies suggest an increased risk of treatment-emergent hyperglycemia-related adverse reactions in patients treated with the atypical antipsychotics. While relative risk estimates are inconsistent, the association between atypical antipsychotics and increased glucose levels appears to fall on a continuum and olanzapine appears to have a greater association than some other atypical antipsychotics.

Mean increases in blood glucose have been observed in patients treated (median exposure of 9.2 months) with olanzapine in phase 3 of the Clinical Antipsychotic Trials of Intervention Effectiveness (CATIE). The mean increase of serum glucose (fasting and nonfasting samples) from baseline to the average of the 2 highest serum concentrations was 15.9 mg/dL.

In a study of healthy volunteers, subjects who received olanzapine (N=22) for 3 weeks had a mean increase compared to baseline in fasting blood glucose of 2.3 mg/dL. Placebo-treated subjects (N=19) had a mean increase in fasting blood glucose compared to baseline of 0.34 mg/dL.

In an analysis of 7 controlled clinical studies, 2 of which were placebo-controlled, with treatment duration up to 12 weeks, olanzapine and fluoxetine was associated with a greater mean change in random glucose when compared to placebo (+8.65 mg/dL vs. -3.86 mg/dL). The difference in mean changes between olanzapine and fluoxetine and placebo was greater in patients with evidence of glucose dysregulation at baseline (including those patients diagnosed with diabetes mellitus or related adverse reactions). Patients treated with anti-diabetic agents, patients with a baseline random glucose level ≥200 mg/dL, or a baseline fasting gluc

dystonia have been reported infrequently (<1%) with the olanzapine and fluoxetine combination.

Additional Findings Observed in Clinical Studies

Sexual Dysfunction — In the pool of controlled olanzapine and fluoxetine studies in patients with bipolar depression, there were higher rates of the treatment-emergent adverse reactions decreased libido, anorgasmia, erectile dysfunction and abnormal ejaculation in the olanzapine and fluoxetine group than in the placebo group. One case of decreased libido led to discontinuation in the olanzapine and fluoxetine group. In the controlled studies that contained a fluoxetine arm, the rates of decreased libido and abnormal ejaculation in the olanzapine and fluoxetine group were less than the rates in the fluoxetine group. None of the differences were statistically significant.

Sexual dysfunction, including priapism, has been reported with all SSRIs. While it is difficult to know the precise risk of sexual dysfunction associated with the use of SSRIs, physicians should routinely inquire about such possible side effects.

There are no adequate and well-controlled studies examining sexual dysfunction with olanzapine and fluoxetine or fluoxetine treatment. Symptoms of sexual dysfunction occasionally persist after discontinuation of fluoxetine treatment.

Difference Among Dose Levels Observed in Other Olanzapine Clinical Trials

In a single 6-week randomized, double-blind, fixed-dose study comparing 5 mg (N=199), 20 (N=200), and 40 (N=200) mg/day of olanzapine in patients with Schizophrenia or Schizoaffective disorder, statistically significant differences among 3 dose groups were observed for the following safety outcomes: weight gain, prolactin elevation, fatigue, and dizziness. Mean baseline to endpoint increase in weight (10 mg/day: 1.9 kg; 20 mg/day: 2.3 kg; 40 mg/day: 3 kg) was observed with significant differences between 10 to 40 mg/day. Increased incidence of prolactin elevation (prolactin >24.2 ng/ml, (female) or >18.77 ng/ml, (male) at any time during the trial (10 mg/day: 31.2%; 20 mg/day: 42.7%; 40 mg/day: 61.1%) with significant differences between 10 vs 40 mg/day and 20 vs 40 mg/day; fatigue (10 mg/day: 3.5%; 20 mg/day: 2.1%; 40 mg/day: 6.6%) with significant differences 2 vs 40 mg/day and 10 vs 40 mg/day; and dizziness (10 mg/day: 2.6%; 20 mg/day: 1.6%; 40 mg/day: 6.6%) with significant differences between 20 vs 40 mg, was observed.

Other Adverse Reactions Observed in Clinical Studies

Following is a list of treatment-emergent adverse reactions reported by patients treated with olanzapine and fluoxetine in clinical trials. This listing is not intended to include reactions (1) already listed in previous tables or elsewhere in labeling, (2) for which a drug cause was remote, (3) which were so general as to be uninformative, (4) which were not considered to have significant clinical importance, or (5) which occurred at a rate equal to or less than placebo.

Reactions are classified by body system using the following definitions: frequent adverse reactions are those occurring in at least 1/100 patients; infrequent adverse reactions are those occurring in 1/100 to 1/1000 patients; and rare reactions are those occurring in fewer than 1/1000 patients.

Body as a Whole — *Frequent:* chills, neck rigidity, photosensitivity reaction; *Rare:* death.[1]

Cardiovascular System — *Frequent:* vasodilatation

Digestive System — *Frequent:* diarrhea, dyspepsia, gastritis, gastroenteritis, nausea and vomiting, peptic ulcer; *Rare:* gastrointestinal hemorrhage, intestinal obstruction, liver fatty degeneration.

Hemic and Lymphatic System — *Frequent:* ecchymosis; *Infrequent:* anemia, thrombocytopenia; *Rare:* leukopenia, purpura.

Metabolic and Nutritional — *Frequent:* generalized edema, weight loss; *Rare:* bilirubinemia, creatine increased, gout.

Musculoskeletal System — *Rare:* osteoporosis.

Nervous System — *Frequent:* amnesia, hypertonia, incoordinated ataxia, buccoglossal syndrome, coma, depersonalization, dysarthria, emotional lability, euphoria, hypokinesia, movement disorder, mycoclonus; *Rare:* hyperkinesia, libido increased, withdrawal syndrome.

Respiratory System — *Infrequent:* epistaxis, yawn; *Rare:* laryngismus.

Skin and Appendages — *Infrequent:* alopecia, dry skin, pruritus; *Rare:* exfoliative dermatitis

Special Senses — *Frequent:* taste disturbance; *Infrequent:* abnormal vision, accommodation, dry eyes.

Urogenital System — *Frequent:* breast pain, menorrhagia[2], urinary frequency, urinary incontinence; *Infrequent:* amenorrhea[2], female dysfunction[3], hypomenorrhea[2], metrorrhagia[2], urinary retention, urinary urgency, urination impaired; *Rare:* breast engorgement[2]

[1] This term represents a serious adverse event but does not meet the definition for adverse drug reactions. It is included here because of its seriousness.

[2] Adjusted for gender.

[3] Adjusted for menopausal status.

Other Adverse Reactions Observed with Olanzapine or Fluoxetine Monotherapy

The following adverse reactions were not observed in olanzapine and fluoxetine-treated patients during premarketing clinical studies but have been reported with olanzapine or fluoxetine monotherapy: aplastic anemia, bruxism, cholelithiasis, cystitis, diabetic coma, eosinophilic pneumonia, exacerbation of multiple sclerosis, gastric ulcer, gynecological bleeding, headache, hyponatremia, jaundice, neutropenia, restless legs syndrome, stuttering[1], sudden unexplained death[1], sweating, and vaginitis. Rechallenge was positive for a few cases of >1000 mg/d, have been reported.

[1] Nothing was only studied in oral and long acting injection (LAI) olanzapine formulations

* These terms represent serious adverse events but do not meet the definition for adverse drug reactions. They are included here because of their seriousness.

Women and Adolescent Patients (ages 10 to 17 years) with a Diagnosis of Bipolar Depression

The information below is derived from a single, 8-week, randomized, placebo-controlled clinical trial investigating olanzapine and fluoxetine for the treatment of bipolar I depression in patients 10 to 17 years of age.

Adverse Reactions Associated with Discontinuation of Treatment in the single pediatric study — Overall, 14.1% of the 170 patients in the olanzapine and fluoxetine group discontinued due to adverse reactions compared with 5.9% of the 85 patients for placebo. Adverse reactions leading to discontinuation associated with the use of olanzapine and fluoxetine (incidence of at least 1% for olanzapine and fluoxetine and greater than that for placebo) using MedDRA Dictionary coding were weight increased (2.9%), suicidal ideation (1.2%), hypersomnia (1.2%), and somnolence (1.2%) versus placebo patients which had 0% incidence of weight increased, bipolar disorder, and somnolence and 1.2% incidence of suicidal ideation.

Adverse Reactions Occurring at an Incidence of 2% or more and greater than placebo — **Table 17** enumerates the treatment-emergent adverse reactions associated with the use of olanzapine and fluoxetine (incidence of at least 2% for olanzapine and fluoxetine and twice or more than placebo).

Table 17: Treatment-Emergent Adverse Reactions: Incidence in a 8-week randomized, double-blind, placebo-controlled clinical trial in pediatric bipolar I depression

System Organ Class	Adverse Reaction	Percentage of Patients Reporting Event	
		Olanzapine and Fluoxetine (N=170)	Placebo (N=85)
Nervous system disorders	Somnolence*	24	2
	Tremor	9	1
Investigations	Weight Increased	20	1
	Blood triglycerides increased	7	2
	Blood cholesterol increased	4	0
	Hepatic Enzyme increased*	3	1
Gastrointestinal disorders	Dyspepsia	3	1
Metabolism and nutrition disorders	Increased appetite	17	1
Psychiatric disorders	Anxiety	3	1
	Restlessness	3	1
	Suicidal Ideation	2	0
Musculoskeletal and connective tissue disorders	Back pain	2	1
Injury, poisoning and procedural complications	Accidental overdose	3	1
Reproductive system and breast disorders	Dysmenorrhea	2	0

* Includes somnolence, sedation, and hypersomnia. No lethargy was reported.

† Includes alanine aminotransferase increased, aspartate aminotransferase increased, hepatic enzyme increased, liver function test abnormal, gamma-glutamyltransferase increased, and transaminases increased.

6.2 Vital Signs and Laboratory Studies

Adults:

Vital Signs — Tachycardia, bradycardia, and orthostatic hypotension have occurred in olanzapine and fluoxetine-treated patients *[see Warnings and Precautions (5.11)]*. The mean standing pulse rate of olanzapine and fluoxetine-treated patients small amount *[see Clinical Pharmacology (12.3)]*. Agents that induce CYP1A2 or glucuronyl transferase enzymes, such as omeprazole and rifampin, may cause an increase in olanzapine clearance. The effect of CYP1A2 inhibitors, such as fluvoxamine and some fluoroquinolone antibiotics, on olanzapine and fluoxetine has not been evaluated. Although olanzapine is metabolized by multiple enzyme systems, induction or inhibition of a single enzyme may appreciably alter olanzapine clearance. Therefore, a dosage increase (for induction) or a dosage decrease (for inhibition) may need to be considered with specific drugs.

7.7 Potential for Olanzapine and Fluoxetine to Affect Other Drugs

Pimozide — Concomitant use of olanzapine and fluoxetine and pimozide is contraindicated. Pimozide can prolong the QT interval. Olanzapine and Fluoxetine can increase the level of pimozide through inhibition of CYP2D6. Olanzapine and Fluoxetine can also prolong the QT interval. Clinical studies of pimozide with other antidepressants demonstrate an increase in drug interaction or QTc prolongation. While a specific study with pimozide and olanzapine and fluoxetine has not been conducted, the potential for drug interactions or QTc prolongation warrants restricting the concurrent use of pimozide and olanzapine and fluoxetine *[see Contraindications (4.2), Warnings and Precautions (5.2)] and Drug Interactions (7.6)]*.

Carbamazepine — Patients on stable doses of carbamazepine have developed elevated plasma anticonvulsant concentrations and clinical anticonvulsant toxicity following initiation of concomitant fluoxetine treatment.

Alcohol — The coadministration of ethanol with olanzapine and fluoxetine may potentiate sedation and orthostatic hypotension *[see Drug Interactions (7.6)]*.

Thioridazine — Thioridazine should not be administered with olanzapine and fluoxetine or administered within a minimum of 5 weeks after discontinuation of olanzapine and fluoxetine capsules, because of the risk of significant prolongation *[see Contraindications (4.2), Warnings and Precautions (5.20), and Drug Interactions (7.8)]*.

In a study of 19 healthy male subjects, which included 6 slow and 13 rapid hydroxylators of debrisoquin, a single 25 mg dose of thioridazine administered after 4 weeks of 20 mg/day dosing of fluoxetine produced a 2.4-fold increase in the slow hydroxylators compared with the rapid hydroxylators. The rate of debrisoquin hydroxylation is felt to depend on the level of CYP2D6 isozyme activity. Thus, this study suggests that fluoxetine, like other drugs that inhibit CYP2D6, such as certain SSRIs, including fluoxetine, will produce elevated plasma levels of thioridazine *[see Contraindications (4.2)]*.

Thioridazine administration produces a dose-related prolongation of the QT$_c$ interval, which is associated with serious ventricular arrhythmias, such as torsades de pointes-type arrhythmias and sudden death. This risk is expected to increase with fluoxetine-induced inhibition of thioridazine metabolism *[see Contraindications (4.2)]*.

Due to the risk of serious ventricular arrhythmias and sudden death potentially associated with elevated thioridazine plasma levels, thioridazine should not be administered with fluoxetine or within a minimum of 5 weeks after fluoxetine has been discontinued *[see Contraindications (4.2)]*.

Tricyclic Antidepressants (TCAs) — Single doses of olanzapine did not affect the pharmacokinetics of imipramine or its active metabolite desipramine.

In 2 fluoxetine studies, steady-state plasma levels of imipramine and desipramine have increased >2-to 10-fold when fluoxetine has been administered in combination. This influence may persist for 3 weeks or longer after fluoxetine is discontinued. Thus, the dose of TCA may need to be decreased, and plasma TCA monitoring may be needed to be monitored temporarily when olanzapine and fluoxetine is coadministered or has been recently discontinued *[see Warnings and Precautions (5.6) Clinical Pharmacology (12.3)]*.

Antihypertensive Agents — Because of the potential for olanzapine to induce hypotension, olanzapine and fluoxetine may enhance the effects of certain antihypertensive agents *[see Warnings and Precautions (5.11)]*.

Benzodiazepines — The clearance component of olanzapine and fluoxetine may antagonize the effects of levodopa and dopamine agonists.

Benzodiazepines — Multiple doses of olanzapine did not influence the pharmacokinetics of diazepam and its active metabolite N-desmethyldiazepam.

When concurrently administered with fluoxetine, the half-life of diazepam may be prolonged in some patients *[see Clinical Pharmacology (12.3)]*. Coadministration of fluoxetine with diazepam has resulted in increased diazepam plasma concentrations and in further psychomotor performance decrement due to increased alprazolam levels.

Lithium — Elevation of blood levels of clozapine has been observed in patients receiving concomitant fluoxetine.

Haloperidol — Fluoxetine has been reported to increase haloperidol blood levels. Patients should be carefully monitored when fluoxetine is added to haloperidol.

Phenytoin — Patients on stable doses of phenytoin have developed elevated plasma levels of phenytoin and clinical phenytoin toxicity following initiation of concomitant fluoxetine.

Drugs Metabolized by CYP2D6 — *In vitro* studies utilizing human liver microsomes suggest that olanzapine has little potential to inhibit CYP2D6. *In vivo* studies of olanzapine and CYP2D6 substrates are consistent with this finding. Olanzapine is a weak inhibitor of CYP2D6 and may inhibit the metabolism of drugs that are metabolized by this enzyme.

Fluoxetine inhibits the activity of CYP2D6 and may make individuals with normal CYP2D6 metabolic activity resemble a poor metabolizer. Coadministration of fluoxetine with other drugs that are metabolized by CYP2D6, including certain antidepressants (TCAs), antipsychotics (phenothiazines and most typical neuroleptics), and antiarrhythmics (e.g., propafenone, flecainide, and others) should be approached with caution. Therapy with medications that are predominantly metabolized by the CYP2D6 system and that have a relatively narrow therapeutic index should be initiated at the low end of the range if the patient is receiving fluoxetine concurrently or has taken it in the previous 5 weeks. Fluoxetine is added to the treatment regimen of a patient already receiving a drug metabolized by CYP2D6, the need for a decrease dose of the CYP2D6 drug should be considered and dose titration should be carefully made. Drugs with a narrow therapeutic index represent the greatest concern (including but not limited to, flecainide, propafenone, vinblastine, and TCAs).

Drugs Metabolized by CYP3A — *In vitro* studies utilizing human liver microsomes suggest that olanzapine has little potential to inhibit CYP3A. Thus, olanzapine is unlikely to cause clinically important drug interactions mediated by these enzymes.

In vivo interaction studies involving the coadministration of fluoxetine with single doses of terfenadine (a CYP3A substrate), no increase in plasma terfenadine concentrations occurred with concomitant fluoxetine. In addition, in vitro studies have shown ketoconazole (a potent inhibitor of CYP3A) to be at least 100 times more potent than fluoxetine or norfluoxetine as an inhibitor of the metabolism of several substrates for this enzyme, including terfenadine, cisapride, astemizole, and midazolam. These data indicate that fluoxetine's extent of inhibition of CYP3A activity is not likely to be of clinical significance.

Effect of Olanzapine on Drugs Metabolized by Other CYP Enzymes — *In vitro* studies utilizing human liver microsomes suggest that olanzapine has little potential to inhibit CYP1A2, CYP2C9, and CYP2C19. Thus, olanzapine is unlikely to cause clinically important drug interactions mediated by these enzymes.

Lithium — Multiple dose fluoxetine does not influence the pharmacokinetics of lithium.

There have been reports of both increased and decreased lithium levels when lithium was used concomitantly with fluoxetine. Cases of lithium toxicity and increased serotonergic effects have been reported. Lithium levels should be monitored in patients taking fluoxetine and olanzapine and fluoxetine concurrently with other *Warnings and Precautions (5.5)]*.

Drugs Tightly Bound to Plasma Proteins — The *in vitro* binding of olanzapine and fluoxetine to human plasma proteins is similar to the individual components. The interaction between olanzapine and fluoxetine and other highly protein-bound drugs has not been fully evaluated. Because fluoxetine is tightly bound to plasma protein, the administration of fluoxetine to a patient taking another drug that is tightly bound to protein (e.g., Coumadin, digitoxin) may cause a shift in plasma concentrations potentially resulting in an adverse effect. Conversely, displacement of bound drug by fluoxetine may result from displacement of protein-bound fluoxetine by other tightly bound drugs *[see Clinical Pharmacology (12.3)]*.

Valproate — *In vitro* studies using human liver microsomes determined that olanzapine has little potential to inhibit the major metabolic pathway, glucuronidation, of valproate. Further, olanzapine has little effect on the metabolism of olanzapine *in vitro*. Thus, a clinically significant pharmacokinetic interaction between olanzapine and valproate is unlikely.

Ibipenten — Multiple doses of olanzapine did not influence the pharmacokinetics of ibipenten.

Theophylline — Multiple doses of olanzapine did not affect the pharmacokinetics of theophylline or its metabolites.

7.8 Drugs that Prolong the QT Interval

Do not use olanzapine and fluoxetine in combination with thioridazine or pimozide. Use olanzapine and fluoxetine with caution in patients receiving other drugs that cause QT prolongation. These include specific antipsychotics (e.g., ziprasidone, lloperidone, chlorpromazine, mesoridazine, droperidol); specific antibiotics (e.g., erythromycin, sparfloxacin, moxifloxacin, gatifloxacin); Class IA antiarrhythmic medications (e.g., quinidine, procainamide); Class III antiarrhythmics (e.g., amiodarone, sotalol); and others (e.g., pentamidine, levomethadyl acetate, methadone, halofantrine, mefloquine, dolasetron mesylate, probucol or tacrolimus). Fluoxetine is primarily metabolized by CYP2D6, a common drug metabolizing enzyme. Coadministration with CYP2D6 inhibitors can increase the concentration of olanzapine and fluoxetine. Consider use of other highly protein-bound drugs can increase the concentration of fluoxetine *[see Contraindications (4.2), Warnings and Precautions (5.20), Drug Interactions (7.7), and Clinical Pharmacology (12.3)]*.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Pregnancy Category C

Risk Summary

There are no adequate and well-controlled clinical studies with olanzapine and fluoxetine or its components (olanzapine and fluoxetine) in pregnant women.

A number of published epidemiological studies assessing the risk of fluoxetine exposure during the first trimester of pregnancy and neonatal outcomes have been reported. Some of these studies have reported an increased risk of congenital malformations, overall, specifically cardiovascular malformations, and premature birth. Other studies have not found an increased risk of congenital malformations overall or cardiovascular malformations specifically. Some studies have reported an increased risk of spontaneous abortion. Results of a number of published epidemiological studies assessing the risk of fluoxetine exposure during the third trimester of pregnancy and neonatal outcomes have been mixed. More than 10 cohort studies and case-control studies failed to demonstrate an increased risk for congenital malformations overall. However, one prospective cohort study conducted by the European Network of Teratology Information Services reported an increased risk of cardiovascular malformations in infants born to women (N = 253) exposed to fluoxetine during the first trimester of pregnancy compared to infants of women (N = 1,340) who were not exposed to fluoxetine. There was no specific pattern of cardiovascular malformations. Overall, however, a causal relationship has not been established.

Neonates exposed to fluoxetine, a component of olanzapine and fluoxetine capsules, and other SSRIs or serotonin and norepinephrine reuptake inhibitors, during late pregnancy have developed complications requiring prolonged hospitalization, respiratory support, and tube feeding. These complications can arise immediately upon delivery. Reported clinical findings have included respiratory distress, cyanosis, apnea, seizures, temperature instability, feeding difficulties, vomiting, hypoglycemia, hypotonia, hypertonia, hyperreflexia, tremor, jitteriness, irritability, and constant crying. These features are consistent with either a direct toxic effect of SSRIs and SNRIs or, possibly, a drug discontinuation syndrome. In some cases, the clinical picture is consistent with serotonin syndrome. Neonates exposed to olanzapine, a component of olanzapine and fluoxetine capsules, during third trimester of pregnancy are at risk for extrapyramidal and/or withdrawal symptoms following delivery including agitation, hypertonia, hypotonia, tremor, somnolence, respiratory distress and feeding disorder. These complications have varied in severity; while in some cases symptoms have been self-limited, in other cases neonates have required intensive care unit support and prolonged hospitalization.

Infants exposed to SSRIs in pregnancy may have an increased risk for persistent pulmonary hypertension of the newborn (PPHN).

Olanzapine and Fluoxetine should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus, taking into account the risk of untreated Bipolar I Depression or Treatment Resistant Depression.

Human Data

There are no adequate and well-controlled clinical studies with olanzapine and fluoxetine, olanzapine or fluoxetine in pregnant women. Olanzapine exposure during pregnancy: data from premarketing clinical studies with olanzapine, including 2 resulting in normal births, 1 resulting in neonatal death due to a cardiovascular defect, 3 therapeutic abortions, and 1 spontaneous abortion. Results of a number of published epidemiological studies assessing the risk of fluoxetine exposure during the first trimester of pregnancy and neonatal outcomes have been mixed. More than 10 cohort studies and case-control studies failed to demonstrate an increased risk for congenital malformations overall. However, one prospective cohort study conducted by the European Network of Teratology Information Services reported an increased risk of cardiovascular malformations in infants born to women (N = 253) exposed to fluoxetine during the first trimester of pregnancy compared to infants of women (N = 1,340) who were not exposed to fluoxetine. There was no specific pattern of cardiovascular malformations. Overall, however, a causal relationship has not been established.

Neonates exposed to fluoxetine, a component of olanzapine and fluoxetine capsules, and other SSRIs or serotonin and norepinephrine reuptake inhibitors, during late pregnancy have developed complications requiring prolonged hospitalization, respiratory support, and tube feeding. Such complications can arise immediately upon delivery. Reported clinical findings have included respiratory distress, cyanosis, apnea, seizures, temperature instability, feeding difficulties, vomiting, hypoglycemia, hypotonia, hypertonia, hyperreflexia, tremor, jitteriness, irritability, and constant crying. These features are consistent with either a direct toxic effect of SSRIs and SNRIs or, possibly, a drug discontinuation syndrome. It should be noted that in some cases the clinical picture is consistent with serotonin syndrome *[see Warnings and Precautions (5.23) and Warnings and Precautions (5.6)]*.

Infants exposed to SSRIs in pregnancy may have an increased risk for persistent pulmonary hypertension of the newborn (PPHN). PPHN occurs in 1 to 2 per 1,000 live births in the general population and is associated with substantial neonatal mortality and morbidity. Several recent epidemiologic studies suggest a positive statistical association between SSRI (including fluoxetine) use in pregnancy and PPHN. Other studies do not show a significant statistical association.

Neonates exposed to antipsychotic drugs (including olanzapine, a component of olanzapine and fluoxetine) during the third trimester of pregnancy are at risk for extrapyramidal and/or withdrawal symptoms following delivery. There have been reports of agitation, hypertonia, hypotonia, tremor, somnolence, respiratory distress and feeding disorder in these neonates. These complications have varied in severity; while in some cases symptoms have been self-limited, in other cases neonates have required intensive care unit support and prolonged hospitalization.

Physicians should also note the results of a prospective longitudinal study of 201 pregnant women with a history of major depression, most of whom were taking antidepressants during pregnancy, in which women who discontinued antidepressant medication during pregnancy showed a significant increase in relapse of their major depression compared to those women who remained on antidepressant medication throughout pregnancy.

When treating a pregnant woman with olanzapine and fluoxetine, the physician should carefully consider both the potential risks of taking an SSRI, along with the potential benefits of treatment (see Warnings and Precautions). This decision can only be made on a case by case basis *[see Dosage and Administration (2.3)]*.

Olanzapine and Fluoxetine should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

Animal Data

Olanzapine and Fluoxetine — Embryo fetal development studies were conducted in rats and rabbits with olanzapine and fluoxetine in low-dose and high-dose combinations. In rats, the doses were; 2 and 4 mg/kg/day (low-dose) and 5 and 10 mg/kg/day (high-dose) administered during organogenesis. The high-dose (10 mg/kg/day) was approximately 0.6 and 0.5 times the MRHD on a mg/m^2 body surface area), and 4 and 8 mg/kg/day (high-dose) [2 and 1 times the MRHD on a mg/m^2 body surface area], and 8 and 8 mg/kg/day (high-dose) [4 and 2 times the MRHD on a mg/m^2 basis, respectively], and 8 and 8 mg/kg/day (high-dose) [8 and 2 times the MRHD on a mg/m^2 basis, respectively], in the rat, 8 and 8 mg/kg/day, respectively, in the rabbit). In the rabbit, there was no evidence of teratogenicity; however, the high-dose combination produced decreased fetal weight and retarded skeletal ossification in conjunction with maternal toxicity. Similarly, in the rat there was no evidence of teratogenicity; however, a decrease in fetal weight was observed with the high-dose combination.

In a pre- and postnatal study conducted in rats, olanzapine and fluoxetine were orally administered during pregnancy and throughout lactation in combination at doses that levels up to 2 (olanzapine) plus 4 (fluoxetine) mg/kg/day [1 and 0.5 times the MRHD on a mg/m^2 body surface area. An elevation of early postnatal mortality (survival through postnatal day 4 was 9% per litter) and reduced bodyweight (approximately 8% in female) occurred among offspring at the highest dose; the no-effect dose was 0.5 (olanzapine) plus 1 (fluoxetine) mg/kg/day (0.25 and 0.12 times the MRHD on a mg/m^2 body surface area). Among the surviving progeny, there were no adverse effects on physical or neurobehavioral development and reproductive performance at any dose.

Olanzapine — In oral embryo-fetal studies in rats and rabbits, there was no evidence of teratogenicity following administration of olanzapine at doses up to 18 and 30 mg/kg/day, respectively (9 and 30 times the MRHD on a mg/m^2 basis, respectively). In a rat oral pre-/postnatal study, reductions in pup weight and increased pup mortality were observed at a dose of 18 mg/kg/day (9 times the MRHD on a mg/m^2 basis). Gestation was prolonged at 10 mg/kg/day (5 times the MRHD on a mg/m^2 basis). Olanzapine was administered to female rats before mating and during gestation and decreased fetal weight occurred at a maternally toxic dose of 30 mg/kg/day (30 times the MRHD on a mg/m^2 basis). Olanzapine crossed the placenta and was detected in fetal tissues following oral administration. Fetal concentrations were 50% of those in maternal plasma.

Fluoxetine — Fluoxetine was administered orally to pregnant rats and rabbits, there was no evidence of teratogenicity following administration of fluoxetine at doses up to 12.5 and 15 mg/kg/day, respectively (1.5 and 3.6 times, respectively, the MRHD of 80 mg/day on a mg/m^2 basis) throughout organogenesis. However, in rat reproduction studies, an increase in stillborn pups, a decrease in pup weight, and an increase in pup deaths during the first 7 days postpartum occurred following maternal exposure to 12 mg/kg/day (1.5 times the MRHD on a mg/m^2 basis) during gestation or 7.5 mg/kg/day (0.9 times the MRHD on a mg/m^2 basis) during gestation and lactation. There was no evidence of developmental neurotoxicity in offspring of rats exposed to fluoxetine during gestation. The no-effect dose for rat pup mortality was 5 mg/kg/day (0.6 times the MRHD on a mg/m^2 basis).

8.2 Labor and Delivery

Olanzapine and Fluoxetine — The effect of olanzapine and fluoxetine on labor and delivery in humans is unknown. Parturition in rats was not affected by olanzapine and fluoxetine. Olanzapine and Fluoxetine should be used during labor and delivery only if the potential benefit justifies the potential risk.

Olanzapine — The effect of olanzapine on labor and delivery in humans is unknown. Parturition in rats was not affected by olanzapine.

Fluoxetine — The effect of fluoxetine on labor and delivery in humans is unknown. Fluoxetine crosses the placenta; therefore, there is a possibility that fluoxetine may be associated with adverse effects on the newborn.

8.3 Nursing Mothers

Olanzapine and Fluoxetine — Studies evaluating the individual components of olanzapine and fluoxetine in nursing mothers are described below. Because of the potential for serious adverse reactions in nursing infants from olanzapine and fluoxetine, a decision should be made whether to discontinue nursing or to discontinue the drug, taking into account the importance of the drug to the mother. It is recommended that women not breast-feed when taking Olanzapine and Fluoxetine.

Olanzapine — In a study in lactating, healthy women, olanzapine was excreted in breast milk. Mean infant dose at steady-state was estimated to be 1.8% of the maternal olanzapine dose. It is recommended that women receiving olanzapine should not breast-feed.

Fluoxetine — Fluoxetine is excreted in human breast milk. In 1 breast milk sample, the concentration of fluoxetine plus norfluoxetine was 70.4 ng/mL. The concentration in the mother's plasma was 295.0 ng/mL. No adverse effects on the infant were reported. In another case, an infant nursed by a mother on fluoxetine developed crying, sleep disturbance, vomiting, and watery stools. The infant's plasma drug levels were 340 ng/mL of fluoxetine and 208mL of norfluoxetine on the 2nd day of feeding.

8.4 Pediatric Use

Olanzapine and Fluoxetine

The safety and efficacy of olanzapine and fluoxetine in patients 10 to 17 years of age has been established for the acute treatment of Depressive Episodes Associated with Bipolar I Disorder in a single 8-week randomized, placebo-controlled clinical trial *[see Clinical Studies (14.1)]*. Patients were initiated at a dose of 3/25 mg/day and dose-forced titrated to the maximum dose of 12/50 mg/day over two weeks. After Week 2, there was flexible dosing of olanzapine and fluoxetine in the range of 6/25, 6/50, 12/25, or 12/50 mg/day. The average dose was 9.3/37.7 mg and flexible dosing of olanzapine was 6/25 mg/day (lower than the adult). Flexible dosing is recommended, rather than the forced titration used in the study *[see Dosage and Administration (2.1)]*.

The types of adverse events observed with olanzapine and fluoxetine in children and adolescents were generally similar to those observed in adults. However, the magnitude of the increase in weight was greater in children and adolescents than adults. These included increases in lipids, hepatic enzymes, and prolactin, as well as orthostatic hypotension *[see Warnings and Precautions (5.2, 5.20, 5.22), and Vital Signs and Laboratory Studies (6.2)]*. The frequency of weight gain ≥7%, and the magnitude and frequency of increases in lipids, hepatic analytes, and prolactin in children and adolescents treated with olanzapine and fluoxetine were similar to those observed in adolescents treated with olanzapine monotherapy.

The safety and efficacy of olanzapine and fluoxetine in the treatment of bipolar I depression in patients under the age of 10 years have not been established. The safety and effectiveness of olanzapine and fluoxetine in combination for treatment resistant depression in patients less than 18 years of age have not been established.

Anyone considering the use of olanzapine and fluoxetine in a child or adolescent must balance the potential risks with the clinical need *[see Boxed Warning and Warnings and Precautions (5.1)]*.

Olanzapine — Safety and effectiveness in pediatric patients less than 13 years of age have not been established.

Compared to patients from adult clinical trials, adolescents treated with oral olanzapine were likely to gain more than 7% bodyweight, experience increases in fasting triglycerides, total cholesterol, triglycerides, LDL cholesterol, prolactin and hepatic aminotransferase levels.

Animal Data

Fluoxetine — Juvenile animal toxicity studies were performed for fluoxetine alone. Significant toxicity on muscle tissue, neurobehavior, reproductive organs, and bone development has been observed following exposure of juvenile rats to fluoxetine. Administration of fluoxetine (10 mg/kg/day) to rats from weaning to maturity (postnatal day 21 through adulthood day 90 at 3, 10, or 30 mg/kg/day was associated with testicular degeneration and necrosis, decreased epididymal weight and hypertrophy of the prostate gland, and impaired reproductive performance. In approximately 5 to 10 times the average AUC in pediatric patients at the MRHD of 20 mg/day), increased serum levels of creatine phosphokinase, osteopenia, and bone fragility in females, and an increase in the incidence of urinary bladder dilatation (20 mg/day), skeletal muscle degeneration and necrosis, decreased femur length/growth and body weight gain (at 5 to 10 times the average AUC in pediatric patients at the MRHD of 20 mg/day). The no-effect dose for rat toxicity exceeded a maximum tolerated dose. When animals were evaluated after a drug-free period (up to 11 weeks after cessation of dosing), fluoxetine-induced effects on musculoskeletal and reproductive parameters persisted. At a dose of 3 mg/kg/day (approximately 0.1 to 0.2 times the average AUC in pediatric patients at the MRHD and learning deficit at the high dose), and reproductive functional impairment (decreased mating at all doses and impaired fertility at the high dose level). In addition, the testicular and epididymal microscopic lesions and/or decreased sperm concentrations in the mid and high dose group were also observed, indicating that the drug effects on reproductive organs are irreversible. The significance of these findings for humans is unknown.

These fluoxetine toxicities in juvenile rats have not been observed in adult animals. Plasma exposures (AUC) associated with these effects were 0.1, 0.7, 1.4, and 3.5 times the AUC in pediatric patients at the MRHD of 20 mg/day given 1 to 2, and 5 to 10 times, respectively, the average exposure in pediatric patients receiving the MRHD of 20 mg/day. The clinical significance of the findings for metabolite, norfluoxetine, are approximately 0.3 to 0.8, 1 to 8, and 3 to 20 times, respectively, the pediatric exposure at the MRHD.

A specific effect on bone development was reported in juvenile mice administered fluoxetine by the intraperitoneal route to a dose that was 0.03 times the MRHD dose of 20 mg/day on a mg/m^2 basis. A decrease of 20 mg/gm/day in bone density was a decrease in bone mineralization and density at both doses, but the overall growth (body weight gain or femur length) was not affected.

8.5 Geriatric Use

Olanzapine and Fluoxetine — Clinical studies of olanzapine and fluoxetine did not include sufficient numbers of patients ≥65 years of age to determine whether they respond differently from younger patients. Other reported clinical experience has not identified differences in responses between the elderly and younger patients. In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy *[see Dosage and Administration (2.3)]*.

Olanzapine — Of the 2500 patients in premarketing clinical studies with olanzapine, 11% (263 patients) were ≥65 years of age. In addition, several hundred Schizophrenia patients who participated in clinical studies were elderly and in many cases, had concomitant medical illnesses. No overall differences in safety or effectiveness were observed between these subjects and younger subjects, and other reported clinical experience has not identified differences in responses between the elderly and younger patients, but greater sensitivity of some older individuals cannot be ruled out. SNRIs and SSRIs, including olanzapine and fluoxetine, have been associated with cases of clinically significant hyponatremia in elderly patients, who may be at greater risk for this adverse reaction *[see Warnings and Precautions (5.17)]*.

8.6 Hepatic Impairment

In subjects with cirrhosis of the liver, the clearances of fluoxetine and its active metabolite, norfluoxetine, were decreased, thus increasing the elimination half-lives of these substances. A lower or less frequent dose of the fluoxetine-component may be appropriate for patients with cirrhosis of the liver. Caution should be used when titrating olanzapine and fluoxetine capsules in patients with diseases or conditions that could affect its metabolism *[see Dosage and Administration (2.3) and Clinical Pharmacology (12.4)]*.

9 DRUG ABUSE AND DEPENDENCE

9.3 Dependence

Olanzapine and Fluoxetine, as with fluoxetine and olanzapine, has not been systematically studied in humans for its potential for abuse, tolerance, or physical dependence. While the clinical studies did not reveal any tendency for any of these behaviors, these observations were not systematic and it is not possible to predict on the basis of this limited experience the extent to which a CNS-active drug will be misused, diverted, and/or abused once marketed. Consequently, physicians should carefully evaluate patients for history of drug abuse and follow such patients closely, observing them for signs of misuse or abuse of olanzapine and fluoxetine (e.g., development of tolerance, incrementation of dose, drug-seeking behavior).

In studies in rats and rhesus monkeys designed to assess abuse and dependence potential, olanzapine alone was shown to have acute dependence QNS effects similar to, but milder or potentially less severe than, those of pentobarbital at doses up to 15 (rat) and 5 (monkey) times the MRHD (20 mg) on a mg/m^2 basis.

10 OVERDOSAGE

Olanzapine and Fluoxetine — During premarketing clinical studies of olanzapine and fluoxetine in combination, overdose of both fluoxetine and olanzapine were reported in 5 study subjects. Four of the 5 subjects experienced loss of consciousness (3) or coma (1). No fatalities occurred.

Adverse reactions reported with overdose of olanzapine in combination, and olanzapine and fluoxetine, have been reported. An overdose of combination therapy is defined as confirmed or suspected ingestion of a dose of >20 mg olanzapine in combination with a dose of >80 mg fluoxetine. Adverse reactions associated with these reports included somnolence (sedation), impaired coordination (ataxia), impaired neurologic function (talkin, confusion, convulsions, dysarthria), arrhythmias, lethargy, essential tremor, agitation, acute psychosis, hypotension, hypertension, and aggression. Fatalities have been corroborated by exposure to additional substances including alcohol, thioridazine, oxycodone, and propoxyphene.

Overdose of olanzapine and fluoxetine capsules with olanzapine alone, symptoms have been reported in the majority of cases. In symptomatic patients, symptoms with ≥10% incidence included agitation/aggressiveness, dysarthria, speech disorder, tachycardia, reduced level of consciousness ranging from somnolence to deep sedation to coma. Among less commonly reported symptoms were the following potentially medically serious reactions: delirium, grand mal convulsions, cardiac arrhythmias (such as supraventricular tachycardia as well as a patient that experienced sinus pause with spontaneous res