HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use OLANZAPINE and FLUOXETINE CAPSULES, safely and effectively. See full prescribing information for OLANZAPINE and FLUOXETINE CAPSULES.

OLANZAPINE and FLUOXETINE CAPSULES, for oral use

Initial U.S. Approval: 2003

WARNING: SUICIDAL THOUGHTS AND BEHAVIORS; AND INCREASED MORTALITY IN ELDERLY PATIENTS WITH DEMENTIA-RELATED PSYCHOSIS

See full prescribing information for complete boxed warning.

- **Increased risk of suicidal thinking and behavior in children, adolescents, and young adults taking antidepressants. Olanzapine and Fluoxetine capsules are not approved for use in children less than 10 years of age (5.1, 8.4, 17.2).**
- **Monitor for worsening and emergence of suicidal thoughts and behaviors (5.1, 17.2).**
- **Elderly patients with dementia-related psychosis treated with antipsychotic drugs are at an increased risk of death. Olanzapine and Fluoxetine Capsules are not approved for the treatment of patients with dementia-related psychosis (5.2, 5.21, 17.3).**

RECENT MAJOR CHANGES

Warnings and Precautions	
Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS)	(5.4) 10/2016
Serotonin Syndrome	(5.6) 01/2017
Falls	(5.12) 02/2017

INDICATIONS AND USAGE

Olanzapine and Fluoxetine Capsules combines olanzapine, an atypical antipsychotic and fluoxetine, a selective serotonin reuptake inhibitor, indicated for treatment of:

- Acute Depressive Episodes Associated with Bipolar I Disorder (**1**)
- Treatment Resistant Depression (**1**)

DOSAGE AND ADMINISTRATION

- Adult Starting Dose: 6 mg olanzapine with 25 mg fluoxetine (6 mg/25 mg, once daily in the evening (**2.1, 2.2**).
- Adult Maximum Dose: 12 mg/50 mg once daily (**2.1, 2.2**).
- Pediatric Bipolar Depression Starting Dose: 3 mg/25 mg once daily (for ages 10 to 17 years) (**2.1**).
- Pediatric Bipolar Depression Maximum Dose: 12 mg/50 mg (**2.1**).
- Starting dose in patients predisposed to hypotensive reactions, hepatic impairment, or with potential for slowed metabolism; 3 mg/25 mg to 6 mg/25 mg. Escalate dose cautiously (**2.3**)

DOSAGE FORMS AND STRENGTHS

Capsules: 3 mg/25 mg, 6 mg/25 mg, 6 mg/50 mg, 12 mg/25 mg, and 12 mg/50 mg olanzapine/fluoxetine (equivalent to fluoxetine) (**3**)

CONTRAINDICATIONS

- Monoamine Oxidiase Inhibitors (MAOI): Because of the risk of serotonin syndrome, do not use MAOIs intended to treat psychiatric disorders with Olanzapine and Fluoxetine Capsules or within 5 weeks of stopping treatment with Olanzapine and Fluoxetine Capsules. Do not start Olanzapine and Fluoxetine Capsules for at least 14 days of stopping an MAOI intended to treat psychiatric disorders. In addition, do not start Olanzapine and Fluoxetine Capsules in a patient who is being treated with linezolid or intravenous methylene blue. (**4.1**)
- Pimozide: Do not use. Risk of QT prolongation (**4.2, 5.20, 7.7, 7.8**)
- Thioridazine: Do not use. Risk of QT interval prolongation. Do not use thioridazine within 5 weeks of discontinuing olanzapine and fluoxetine capsules (**4.2, 5.20, 7.7, 7.8**)

WARNINGS AND PRECAUTIONS

- *Neuroleptic Malignant Syndrome*: Manage with immediate discontinuation and close monitoring (**5.3**)
- *Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS)*: Discontinue if DRESS is suspected (**5.4**).
- *Metabolic Changes*: Atypical antipsychotic drugs have been associated with metabolic changes including hyperglycemia, dyslipidemia, and weight gain (**5.5**)
  - *Hyperglycemia and Diabetes Mellitus*: In some cases extreme and associated with ketoacidosis or hyperosmolar coma or death. Monitor for symptoms of hyperglycemia. Perform fasting blood glucose testing before beginning, and periodically during treatment. (**5.5**)
  - *Dyslipidemia*: Appropriate clinical monitoring is recommended, including fasting blood lipid testing before beginning, and periodically during, treatment (**5.5**)
  - *Weight Gain*: Consider potential consequences of weight gain. Monitor weight regularly (**5.5**)
- *Serotonin Syndrome*: Serotonin syndrome has been reported with SSRIs and SNRIs, including Olanzapine and Fluoxetine Capsules, both when taken alone, but especially when coadministered with other serotonergic agents (triptans, tricyclic antidepressants, fentanyl, lithium, tramadol, tryptophan, buspirone and St. John's Wort). If such symptoms occur, discontinue Olanzapine and Fluoxetine Capsules, and initiate supportive treatment. If concomitant use of Olanzapine and Fluoxetine Capsules with other serotonergic drugs is clinically warranted, patients should be made aware of the potential increased risk for serotonin syndrome, particularly during treatment initiation and dose increases (**5.6**).
- *Angle-Closure Glaucoma*: Angle-closure glaucoma has occurred in patients with untreated anatomically narrow angles treated with antidepressants (**5.7**)
- *Allergic Reactions and Rash*: Discontinue upon appearance of rash or allergic phenomena (**5.8**)
- *Activation of Mania/Hypomania*: Screen for Bipolar Disorder and monitor for activation of mania/hypomania (**5.9**)
- *Tardive Dyskinesia*: Discontinue if clinically appropriate (**5.10**)
- *Orthostatic Hypotension*: Can be associated with bradycardia and syncope. Risk is increased during initial dose titration. Use caution in patients with cardiovascular disease or cerebrovascular disease, and those conditions that could affect hemodynamic responses (**5.11**)
- *Leukopenia, Neutropenia, and Agranulocytosis*: Has been reported with antipsychotics, including Olanzapine and Fluoxetine. Patients with a history of a clinically significant low white blood cell count (WBC) or drug induced leukopenia/neutropenia should have their complete blood count (CBC) monitored frequently during the first few months of therapy. Consider discontinuing Olanzapine and Fluoxetine Capsules at the first sign of a clinically significant decline in WBC in the absence of other causative factors (**5.13**)
- *Seizures*: Use cautiously in patients with a history of seizures or with conditions that lower the seizure threshold (**5.15**)
- *Abnormal Bleeding*: SSRIs increase the risk of bleeding. Use with NSAIDs, aspirin, warfarin, or other drugs that affect coagulation may potentiate the risk of gastrointestinal or other bleeding (**5.16**)
- *Hyponatremia*: Can occur in association with syndrome of inappropriate antidiuretic hormone (SIADH). Consider discontinuing Olanzapine and Fluoxetine capsules if symptomatic hyponatremia occurs (SIADH) (**5.17**)
- *Potential for Cognitive and Motor Impairment*: Has potential to impair judgment, thinking, and motor skills. Caution patients about operating machinery (**5.18**)
- *QT Prolongation*: QT prolongation and ventricular arrhythmia including Torsade de Pointes have been reported with fluoxetine. Use with caution in conditions that predispose to arrhythmias or increased fluoxetine exposure. Use cautiously in patients with risk factors for QT prolongation (**4.2, 5.20**)
- *Hyperprolactinemia*: Like other antipsychotic-controlled trials of antidepressant drugs (SSRIs and others) showed that these drugs increase the risk of suicidal thinking and behavior (suicidality) in children, adolescents, and young adults (ages 18 to 24) with Major Depressive Disorder (MDD) and other psychiatric disorders. Short-term studies did not show an increase in the risk of suicidality with these drugs compared to placebo in adults beyond age 24; there was a reduction with antidepressants compared to placebo in adults aged 65 and older. (**5.19**)
- *Long Elimination Half-Life of Fluoxetine*: Changes in dose will not be fully reflected in plasma for several weeks (**5.24**)

ADVERSE REACTIONS

Most common adverse reactions (≥ 5% and at least twice that for placebo) in adults: sedation, weight increased, appetite increased, dry mouth, fatigue, edema, tremor, disturbance in attention, blurred vision. Children and adolescents: sedation, weight increased, appetite increased, tremor, triglyceride increased, hepatic enzymes increased (**6.1**)

To report SUSPECTED ADVERSE REACTIONS, contact Par Pharmaceutical at 1-800-828-9393 or FDA at 1-800- FDA-1088 or www.fda.gov/medwatch.

DRUG INTERACTIONS

- *Monoamine Oxidase Inhibitor (MAOI)*: (**2.4, 2.5, 4.1, 5.6, 7.1**)
- *Drugs Metabolized by CYP2D6*: Fluoxetine is a potent inhibitor of CYP2D6 enzyme pathway (**7.1**)
- *Tricyclic Antidepressants (TCAs)*: Monitor TCA levels during coadministration with olanzapine and fluoxetine or when olanzapine and fluoxetine capsules have been recently discontinued (**7.1**)
- *CNS Acting Drugs*: Caution is advised if the concomitant administration of olanzapine and fluoxetine and other CNS-active drugs is required (**7.2**)
- *Antihypertensive Agent*: Enhanced antihypertensive effect (**7.7**)
- *Levodopa and Dopamine Agonists*: May antagonize levodopa/dopamine agonists (**7.7**)
- *Benzodiazepines*: May potentiate orthostatic hypotension and sedation (**7.6, 7.7**)
- *Clozapine*: May elevate clozapine levels (**7.7**)
- *Haloperidol*: Elevated haloperidol levels have been observed (**7.7**)
- *Carbamazepine*: Potential for elevated carbamazepine levels and clinical anticonvulsant toxicity (**7.7**)
- *Phenytoin*: Potential for elevated phenytoin levels and clinical anticonvulsant toxicity (**7.7**)
- *Alcohol*: May potentiate sedation and orthostatic hypotension (**7.7**)
- *Serotonergic Drugs*: (**2.4, 2.5, 4.1, 5.6, 7.3**)
- *Fluvoxamine*: May increase olanzapine levels; a lower dose of the olanzapine component of olanzapine and fluoxetine capsules should be considered (**7.8**)
- *Drugs that Interfere with Hemostasis (e.g., NSAIDs, Aspirin, Warfarin, etc.)*: May potentiate the risk of bleeding (**7.4**)
- *Drugs Tightly Bound to Plasma Proteins*: Fluoxetine may cause shift in plasma concentrations (**7.7**)
- *Drugs that Prolong the QT Interval*: Do not use olanzapine and fluoxetine capsules in combination with thioridazine or pimozide. Use olanzapine and fluoxetine capsules with caution in combination with other drugs that prolong the QT interval (**4.2, 5.20, 7.7, 7.8**)

USE IN SPECIFIC POPULATIONS

- *Pregnancy*: Olanzapine and Fluoxetine capsules should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus (**8.1**)
- *Nursing Mothers*: Breast feeding is not recommended (**8.3**)
- *Pediatric Use*: Safety and efficacy of olanzapine and fluoxetine capsules for the treatment of bipolar I depression in patients under 10 years of age have not been established. Safety and efficacy of olanzapine and fluoxetine capsules for treatment resistant depression in patients under 18 years of age have not been established (**8.4**).
- *Hepatic Impairment*: Use a lower or less frequent dose in patients with cirrhosis (**8.6**)

See 17 for PATIENT COUNSELING INFORMATION and the FDA-approved Medication Guide

FULL PRESCRIBING INFORMATION: CONTENTS *

WARNINGS: SUICIDAL THOUGHTS AND BEHAVIORS; AND INCREASED MORTALITY IN ELDERLY PATIENTS WITH DEMENTIA-RELATED PSYCHOSIS

1 INDICATIONS AND USAGE
2 DOSAGE AND ADMINISTRATION
2.1 Depressive Episodes Associated with Bipolar I Disorder
2.2 Treatment Resistant Depression
2.3 Specific Populations
2.4 Switching a Patient To or From a Monoamine Oxidase Inhibitor (MAOI) Intended to Treat Psychiatric Disorders
2.5 Use of Olanzapine and Fluoxetine Capsules with other MAOIs such as Linezolid or Methylene Blue
2.6 Discontinuation of Treatment with Olanzapine and Fluoxetine Capsules
3 DOSAGE FORMS AND STRENGTHS
4 CONTRAINDICATIONS
4.1 Monoamine Oxidase Inhibitors (MAOIs)
4.2 Pimozide and Thioridazine
5 WARNINGS AND PRECAUTIONS
5.1 Suicidal Thoughts and Behaviors in Children, Adolescents, and Young Adults
5.2 Increased Mortality in Elderly Patients with Dementia-Related Psychosis
5.3 Neuroleptic Malignant Syndrome (NMS)
5.4 Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS)
5.5 Metabolic Changes
5.6 Serotonin Syndrome
5.7 Angle-Closure Glaucoma
5.8 Allergic Reactions and Rash
5.9 Activation of Mania/Hypomania
5.10 Tardive Dyskinesia
5.11 Orthostatic Hypotension
5.12 Falls
5.13 Leukopenia, Neutropenia and Agranulocytosis
5.14 Dysphagia
5.15 Seizures
5.16 Abnormal Bleeding
5.17 Hyponatremia
5.18 Potential for Cognitive and Motor Impairment
5.19 Body Temperature Dysregulation
5.20 QT Prolongation
5.21 Use in Patients with Concomitant Illness
5.22 Hyperprolactinemia
5.23 Concomitant Use of Olanzapine and Fluoxetine Products
5.24 Long Elimination Half-Life of Fluoxetine
5.25 Discontinuation Adverse Reactions
6 ADVERSE REACTIONS
6.1 Clinical Trials Experience
6.2 Vital Signs and Laboratory Studies
6.3 Postmarketing Experience
7 DRUG INTERACTIONS
7.1 Monoamine Oxidase Inhibitors (MAOIs)
7.2 CNS Acting Drugs
7.3 Serotonergic Drugs
7.4 Drugs that Interfere with Hemostasis (e.g., NSAIDs, Aspirin, Warfarin)
7.5 Electroconvulsive Therapy (ECT)
7.6 Potential for Other Drugs to Affect Olanzapine and Fluoxetine
7.7 Potential for Olanzapine and Fluoxetine to Affect Other Drugs
7.8 Drugs that Prolong the QT Interval
8 USE IN SPECIFIC POPULATIONS
8.1 Pregnancy
8.2 Labor and Delivery
8.3 Nursing Mothers
8.4 Pediatric Use
8.5 Geriatric Use
8.6 Hepatic Impairment
9 DRUG ABUSE AND DEPENDENCE
9.3 Dependence
10 OVERDOSAGE
10.1 Management of Overdose
11 DESCRIPTION
12 CLINICAL PHARMACOLOGY
12.1 Mechanism of Action
12.2 Pharmacodynamics
12.3 Pharmacokinetics
12.4 Specific Populations
13 NONCLINICAL TOXICOLOGY
13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility
14 CLINICAL STUDIES
14.1 Depressive Episodes Associated with Bipolar I Disorder
14.2 Treatment Resistant Depression
16 HOW SUPPLIED/STORAGE AND HANDLING
16.1 How Supplied
16.2 Storage and Handling
17 PATIENT COUNSELING INFORMATION
17.1 Information on Medication Guide
17.2 Suicidal Thoughts and Behaviors in Children, Adolescents, and Young Adults
17.3 Elderly Patients with Dementia-Related Psychosis: Increased Mortality and Cerebrovascular Adverse Events (CVAE), Including Stroke
17.4 Neuroleptic Malignant Syndrome (NMS)
17.5 Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS)
17.6 Hyperglycemia and Diabetes Mellitus
17.7 Dyslipidemia
17.8 Weight Gain
17.9 Serotonin Syndrome
17.10 Angle-Closure Glaucoma
17.11 Allergic Reactions and Rash
17.12 Orthostatic Hypotension
17.13 Abnormal Bleeding
17.14 Hyponatremia
17.15 Potential for Cognitive and Motor Impairment
17.16 Body Temperature Dysregulation
17.17 Concomitant Medication
17.18 Discontinuation of Treatment with Olanzapine and Fluoxetine Capsules
17.19 Alcohol
17.20 Use in Specific Populations
17.21 QT Prolongation

* Sections or subsections omitted from the full prescribing information are not listed

FULL PRESCRIBING INFORMATION

WARNINGS: SUICIDAL THOUGHTS AND BEHAVIORS; AND INCREASED MORTALITY IN ELDERLY PATIENTS WITH DEMENTIA-RELATED PSYCHOSIS

Suicidal Thoughts and Behaviors — Antidepressants increased the risk of suicidal thoughts and behavior in children, adolescents, and young adults in short-term studies. These studies did not show an increase in the risk of suicidal thoughts and behavior with antidepressant use in patients over age 24; there was a reduction in risk with antidepressant use in patients aged 65 and older.

In patients of all ages who are started on antidepressant therapy, monitor closely for worsening and emergence of suicidal thoughts and behaviors, anxiety, agitation, and other symptoms. Advise patients, their families, and caregivers of the need for close observation and communication with the prescriber. Olanzapine and Fluoxetine Capsules are not approved for use in children less than 10 years of age (see Warnings and Precautions (5.1), Use in Specific Populations (8.4), and Patient Counseling Information (17.2)).

Increased Mortality in Elderly Patients with Dementia-Related Psychosis — Elderly patients with dementia-related psychosis treated with antipsychotic drugs are at an increased risk of death. Olanzapine and Fluoxetine Capsules are not approved for the treatment of patients with dementia-related psychosis (see Warnings and Precautions (5.2)).

1 INDICATIONS AND USAGE

Olanzapine and Fluoxetine is indicated for the treatment of:

- Acute depressive episodes in Bipolar I Disorder (see *Clinical Studies (14.1)*).
- Treatment resistant depression (Major Depressive Disorder in patient who do not respond to 2 adequate trials of different antidepressants of adequate dose and duration in the current episode) (see *Clinical Studies (14.2)*).

2 DOSAGE AND ADMINISTRATION

2.1 Depressive Episodes Associated with Bipolar I Disorder

Adults – Administer Olanzapine and Fluoxetine Capsules once daily in the evening, generally beginning with the 6 mg/25 mg (mg olanzapine/mg equivalent fluoxetine) capsule. While food has no appreciable effect on the absorption of olanzapine and fluoxetine given individually, the effect of food on the absorption of olanzapine and fluoxetine has not been studied. Make dosage adjustments, if indicated, according to efficacy and tolerability. Antidepressant efficacy was demonstrated with olanzapine and fluoxetine in a dose range of olanzapine 6 mg to 12 mg and fluoxetine 25 mg to 50 mg (see *Clinical Studies (14.1)*). The safety of doses above 18 mg of olanzapine and 75 mg of fluoxetine has not been evaluated in adult clinical studies. Periodically reexamine the need for continued pharmacotherapy.

Children and Adolescents (10 to 17 years of age) – Administer Olanzapine and Fluoxetine Capsules once daily in the evening, generally beginning with the 3 mg/25 mg capsule. While food has no appreciable effect on the absorption of olanzapine and fluoxetine given individually, the effect of food on the absorption of olanzapine and fluoxetine has not been studied. Increase dosage, if indicated, according to efficacy and tolerability. Antidepressant efficacy was demonstrated with olanzapine and fluoxetine in a dose range of olanzapine 6 mg to 12 mg and fluoxetine 25 mg to 50 mg (see *Clinical Studies (14.1)*). The safety of doses above 18 mg of olanzapine and 75 mg of fluoxetine has not been evaluated in adult clinical studies. Periodically reexamine the need for continued pharmacotherapy.

2.2 Treatment Resistant Depression

Administer Olanzapine and Fluoxetine Capsules once daily in the evening, generally beginning with the 6 mg/ 25 mg capsule. While food has no appreciable effect on the absorption of olanzapine and fluoxetine given individually, the effect of food on the absorption of Olanzapine and Fluoxetine Capsules has not been studied. Adjust dosage, if indicated, according to efficacy and tolerability.

Antidepressant efficacy was demonstrated with Olanzapine and Fluoxetine Capsules in a dose range of olanzapine 12 mg to 18 mg and fluoxetine 25 mg to 50 mg (see *Clinical Studies (14.2)*). The safety of doses above 18 mg/ 75 mg has not been evaluated in clinical studies. Periodically reexamine the need for continued pharmacotherapy.

2.3 Specific Populations

Start Olanzapine and Fluoxetine Capsules 3 mg/25 mg or 6 mg/25 mg in patients with a predisposition to hypotensive reactions, patients with hepatic impairment, or patients who exhibit a combination of factors that may slow the metabolism of olanzapine and fluoxetine (female gender, geriatric age, nonsmoking status) or those patients who may be pharmacodynamically sensitive to olanzapine. Titrate slowly and adjust dosage as needed in patients who exhibit a combination of factors that may slow metabolism. Olanzapine and fluoxetine capsules were not studied in patients >65 years of age or in patients <10 years of age (see *Warnings and Precautions (5.21), Use in Specific Populations (8.5), and Clinical Pharmacology (12.3, 12.4)*).

Treatment of Pregnant Women – When treating pregnant women with fluoxetine, a component of olanzapine and fluoxetine, the physician should carefully consider the potential risks and potential benefits of treatment. Neonates exposed to SSRIs or SNRIs late in the third trimester have developed complications requiring prolonged hospitalizations, respiratory support, and tube feeding (see *Use in Specific Populations (8.1)*).

2.4 Switching a Patient To or From a Monoamine Oxidase Inhibitor (MAOI) Intended to Treat Psychiatric Disorders

At least 14 days should elapse between discontinuation of an MAOI intended to treat psychiatric disorders and initiation of therapy with olanzapine and fluoxetine. Conversely, at least 5 weeks should be allowed after stopping olanzapine and fluoxetine before starting an MAOI intended to treat psychiatric disorders (see *Contraindications (4.1)*).

2.5 Use of Olanzapine and Fluoxetine Capsules with Other MAOIs such as Linezolid or Methylene Blue

Do not start olanzapine and fluoxetine in a patient who is being treated with linezolid or intravenous methylene blue because there is an increased risk of serotonin syndrome. In a patient who requires more urgent treatment of a psychiatric condition, other interventions, including hospitalization, should be considered (see *Contraindications (4.1)*).

In some cases, a patient already receiving olanzapine and fluoxetine therapy may require urgent treatment with linezolid or intravenous methylene blue. If acceptable alternatives to linezolid or intravenous methylene blue treatment are not available and the potential benefits of linezolid or intravenous methylene blue are judged to outweigh the risks of serotonin syndrome in a particular patient, olanzapine and fluoxetine should be stopped promptly, and linezolid or intravenous methylene blue can be administered. The patient should be monitored for symptoms of serotonin syndrome for five weeks or until 24 hours after the last dose of linezolid or intravenous methylene blue, whichever comes first. Therapy with olanzapine and fluoxetine may be resumed 24 hours after the last dose of linezolid or intravenous methylene blue (see *Warnings and Precautions (5.6)*).

The risk of administering methylene blue by non-intravenous routes (such as oral tablets or by local injection) or in intravenous doses much lower than 1 mg/kg with olanzapine and fluoxetine is unclear. The clinician should, nevertheless, be aware of the possibility of emergence of serotonin syndrome with such use (see *Warnings and Precautions (5.6)*).

2.6 Discontinuation of Treatment with Olanzapine and Fluoxetine Capsules

Symptoms associated with discontinuation of fluoxetine, a component of olanzapine and fluoxetine, SNRIs, and SSRIs, have been reported (see *Warnings and Precautions (5.25)*).

3 DOSAGE FORMS AND STRENGTHS

Capsules (mg olanzapine/mg equivalent fluoxetine):

- 3 mg/25 mg
- 6 mg/25 mg
- 6 mg/50 mg
- 12 mg/25 mg
- 12 mg/50 mg

4 CONTRAINDICATIONS

4.1 Monoamine Oxidase Inhibitors (MAOIs)

The use of MAOIs intended to treat psychiatric disorders with olanzapine and fluoxetine or within 5 weeks of stopping treatment with olanzapine and fluoxetine is contraindicated because of an increased risk of serotonin syndrome. The use of olanzapine and fluoxetine within 14 days of stopping an MAOI intended to treat psychiatric disorders is also contraindicated (see *Dosage and Administration (2.4) and Warnings and Precautions (5.6)*).

Starting olanzapine and fluoxetine in a patient who is being treated with MAOIs such as linezolid or intravenous methylene blue is also contraindicated because of an increased risk of serotonin syndrome *[see Dosage and Administration (2.5) and Warnings and Precautions (5.6)]*.

4.2 Other Contraindications

- Pimozide — *[see Warnings and Precautions (5.20) and Drug Interactions (7.7, 7.8)]*
- Thioridazine — *[see Warnings and Precautions (5.20) and Drug Interactions (7.7, 7.8)]*

Pimozide and thioridazine prolong the QT interval. Olanzapine and Fluoxetine can increase the levels of pimozide and thioridazine through inhibition of CYP2D6. Olanzapine and fluoxetine can also prolong the QT interval.

5 WARNINGS AND PRECAUTIONS

5.1 Suicidal Thoughts and Behaviors in Children, Adolescents, and Young Adults

Patients with Major Depressive Disorder (MDD), both adult and pediatric, may experience worsening of their depression and/or the emergence of suicidal ideation and behavior (suicidality) or unusual changes in behavior, whether or not they are taking antidepressant medications, and this risk may persist until significant remission has occurred. Suicide is a known risk of depression and certain other psychiatric disorders, and these disorders themselves are the strongest predictors of suicide. There has been a long-standing concern, however, that antidepressants may have a role in inducing worsening of depression and the emergence of suicidality in certain patients during the early phases of treatment. Pooled analyses of short-term placebo-controlled trials of antidepressant drugs (SSRIs and others) showed that these drugs increase the risk of suicidal thinking and behavior (suicidality) in children, adolescents, and young adults (ages 18 to 24) with Major Depressive Disorder (MDD) and other psychiatric disorders. Short-term studies did not show an increase in the risk of suicidality with these drugs compared to placebo in adults beyond age 24; there was a reduction with antidepressants compared to placebo in adults aged 65 and older.

The pooled analyses of placebo-controlled trials in children and adolescents with MDD, Obsessive Compulsive Disorder (OCD), or other psychiatric disorders included a total of 24 short-term trials of 9 antidepressant drugs in over 4400 patients. The pooled analyses of placebo-controlled trials in adults with MDD or other psychiatric disorders included a total of 295 short-term trials (median duration of 2 months) of 11 antidepressant drugs in over 77,000 patients. There was considerable variation in risk of suicidality among drugs, but a tendency toward an increase in the younger patients for almost all drug studied. There were differences in absolute risk of suicidality across the different indications, with the highest incidence in MDD. The risk differences (drug versus placebo), however, were relatively stable within age strata and across indications. These risk differences (drug-placebo difference in the number of cases of suicidality per 1000 patients treated) are provided in **Table 1**

Table 1: Suicidality per 1000 Patients Treated

Age Range	Drug-Placebo Difference in Number of Cases of Suicidality per 1000 Patients Treated
	Increases Compared to Placebo
<18	14 additional cases
18 to 24	5 additional cases
	Decreases Compared to Placebo
25 to 64	1 fewer case
≥65	6 fewer cases

No suicides occurred in any of the pediatric trials. There were suicides in the adult trials, but the number was not sufficient to reach any conclusion about drug effect on suicide.

It is unknown whether the susceptibility of children and adolescents to suicidal events, like that observed in adults, would be mitigated by long-term use (i.e., beyond several months. However, there is substantial evidence from placebo-controlled maintenance trials in adults with depression that the use of antidepressants can delay the recurrence of depression.

All patients being treated with antidepressants for any indication should be monitored appropriately and observed closely for clinical worsening, suicidality, and unusual changes in behavior, especially during the initial few months of a course of drug therapy, or at times of dose changes, either increases or decreases.

The following symptoms, anxiety, agitation, panic attacks, insomnia, irritability, hostility, aggressiveness, impulsivity, akathisia (psychomotor restlessness), hypomania, and mania, have been reported in adult and pediatric patients being treated with antidepressants for Major Depressive Disorder as well as for other indications, both psychiatric and nonpsychiatric. Although a causal link between the emergence of such symptoms and either the worsening of depression and/or the emergence of suicidal impulses has not been established, there is concern that such symptoms may represent precursors to emerging suicidality.

Consideration should be given to changing the therapeutic regimen, including possibly discontinuing the medication, in patients whose depression is persistently worse, or who are experiencing emergent suicidality or symptoms that might be precursors to worsening depression or suicidality, especially if these symptoms are severe, abrupt in onset, or were not part of the patient's presenting symptoms.

If the decision has been made to discontinue treatment, medication should be tapered, as rapidly as is feasible, but with recognition that abrupt discontinuation can be associated with certain symptoms *[see Warnings and Precautions (5.25)]*.

Families and caregivers of patients being treated with antidepressants for Major Depressive Disorder or other indications, psychiatric and nonpsychiatric, should be alerted about the need to monitor patients for the emergence of agitation, irritability, unusual changes in behavior, and the other symptoms described above, as well as the emergence of suicidality, and to report such symptoms immediately to healthcare providers. Such monitoring should include daily observation by families and caregivers. Prescriptions for olanzapine and fluoxetine should be written for the smallest quantity of capsules consistent with good patient management, in order to reduce the risk of overdose.

Screening patients for bipolar disorder and olanzapine and fluoxetine capsules are not approved for use in treating any indications in patients less than 10 years of age *[see Use in Specific Populations (8.4)]*.

5.2 Increased Mortality in Elderly Patients with Dementia-Related Psychosis

Elderly patients with dementia-related psychosis treated with antipsychotic drugs are at an increased risk of death. Olanzapine and fluoxetine capsules are not approved for the treatment of patients with dementia-related psychosis *[see Boxed Warning, Warnings and Precautions (5.21), and Patient Counseling Information (17.3)]*

In olanzapine placebo-controlled clinical trials of elderly patients with dementia-related psychosis, the incidence of death in olanzapine-treated patients was significantly greater than placebo-treated patients (3.5% vs 1.5%, respectively).

Meta-Analysis of Antipsychotic Use in Dementia-Related Psychoses — Elderly patients with dementia-related psychosis treated with antipsychotic drugs are at an increased risk of death. Analyses of seventeen placebo-controlled trials (modal duration of 10 weeks), largely in patients taking atypical antipsychotic drugs, revealed a risk of death in drug treated patients of between 1.6 to 1.7 times the risk of death in placebo-treated patients. Over the course of a typical 10 week controlled trial, the rate of death in drug treated patients was about 4.5% compared to a rate of about 2.6% in the placebo group. Although the causes of death were varied, most of the deaths appeared to be either cardiovascular (e.g., heart failure, sudden death) or infectious (e.g., pneumonia) in nature. The observation in these studies suggest that, similar to atypical antipsychotic drugs, treatment with conventional antipsychotic drugs may increase mortality. The extent to which the findings of increased mortality in observational studies may be attributed to the antipsychotic drugs as opposed to some characteristic(s) of the patients is not clear. Olanzapine and Fluoxetine Capsules are not approved for the treatment of patients with dementia-related psychosis *[see Warnings and Precautions (5.21) and Patient Counseling Information (17.3)]*.

Cerebrovascular Adverse Events (CVAE), Including Stroke – Cerebrovascular adverse events (e.g., stroke, transient ischemic attack), including fatalities, were reported in patients in trials of olanzapine in elderly patients with dementia-related psychosis. In placebo-controlled trials, there was a significantly higher proportion of cerebrovascular adverse events in patients treated with olanzapine compared to patients treated with placebo. Olanzapine and olanzapine and fluoxetine capsules are not approved for the treatment of patients with dementia-related psychosis *[see Boxed Warning and Patient Counseling Information (17.3)]*.

5.3 Neuroleptic Malignant Syndrome (NMS)

A potentially fatal symptom complex sometimes referred to as NMS has been reported in association with administration of antipsychotic drugs, including olanzapine. Clinical manifestations of NMS are hyperpyrexia, muscle rigidity, altered mental status, and evidence of autonomic instability (irregular pulse or blood pressure, tachycardia, diaphoresis, and cardiac dysrhythmia). Additional signs may include elevated creatinine phosphokinase, myoglobinuria (rhabdomyolysis), and acute renal failure.

The diagnostic evaluation of patients with this syndrome is complicated. In arriving at a diagnosis, it is important to exclude cases where the clinical presentation includes both serious medical illness (e.g., pneumonia, systemic infection, etc.) and untreated or inadequately treated extrapyramidal signs and symptoms (EPS). Other important considerations in the differential diagnosis include central anticholinergic toxicity, heat stroke, drug fever, and primary central nervous system pathology.

The management of NMS should include: 1) immediate discontinuation of antipsychotic drugs and other drugs not essential to concurrent therapy, 2) intensive symptomatic treatment and medical monitoring, and 3) treatment of any concomitant serious medical problems for which specific treatments are available. There is no general agreement about specific pharmacological treatment regimens for NMS.

If after recovering from NMS, a patient requires treatment with an antipsychotic, the patient should be carefully monitored, since recurrences of NMS have been reported *[see Warnings and Precautions (5.5) and Patient Counseling Information (17.4, 17.9)]*.

5.4 Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS)

Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS) has been reported with olanzapine exposure. DRESS may present with a cutaneous reaction (such as rash or exfoliative dermatitis), eosinophilia, fever, and/or lymphadenopathy, along with systemic complications such as hepatitis, nephritis, pneumonitis, myocarditis, and pericarditis. DRESS is sometimes fatal. Discontinue olanzapine and fluoxetine capsules if DRESS is suspected *[see Patient Counseling Information (17.5)]*.

5.5 Metabolic Changes

Atypical antipsychotic drugs have been associated with metabolic changes including hyperglycemia, dyslipidemia, and weight gain. While all of the drugs in the class have been associated with increased cardiovascular/cerebrovascular risk, olanzapine's specific metabolic profile is presented below.

Hyperglycemia and Diabetes Mellitus

Adults – Physicians should consider the risks and benefits when prescribing olanzapine and fluoxetine capsules to patients with an established diagnosis of diabetes mellitus, or having borderline increased glucose blood level (fasting 100 to 126 mg/dL, random 140 to 200 mg/dL). Patients receiving olanzapine and fluoxetine capsules should be monitored regularly for worsening of glucose control. Patients starting treatment with olanzapine and fluoxetine should undergo fasting blood glucose testing at the beginning of treatment and periodically during treatment. Any patient treated with atypical antipsychotics should be monitored for symptoms of hyperglycemia including polydipsia, polyuria, polyphagia, and weakness. Patients who develop symptoms of hyperglycemia during treatment with atypical antipsychotics should undergo fasting blood glucose testing. In some cases, hyperglycemia has resolved when the atypical antipsychotic was discontinued; however, some patients required continuation of anti-diabetic treatment despite discontinuation of the suspect drug *[see Patient Counseling Information (17.6)]*.

Hyperglycemia, in some cases extreme and associated with ketoacidosis or hyperosmolar coma or death, has been reported in patients treated with atypical antipsychotics, including olanzapine alone, as well as olanzapine taken concomitantly with fluoxetine. Assessment of the relationship between atypical antipsychotic use and glucose abnormalities is complicated by the possibility of an increased background risk of diabetes mellitus in patients with schizophrenia and the increasing incidence of diabetes mellitus in the general population. Epidemiological studies suggest an increased risk of treatment-emergent hyperglycemia-related adverse reactions in patients treated with the atypical antipsychotics. While relative risk estimates are inconsistent, the association between atypical antipsychotics and increases in glucose levels appears to fall on a continuum. Olanzapine appears to have a greater association than some other atypical antipsychotics.

Mean increases in blood glucose have been observed in patients treated (median exposure of 9.2 months) with olanzapine in phase 1 of the Clinical Antipsychotic Trials of Intervention Effectiveness (CATIE). The mean increase of serum glucose (fasting and nonfasting samples) from baseline to the average of the 2 highest serum concentrations was 15.0 mg/dL.

In a study of healthy volunteers, subjects who received olanzapine (N=22) for 3 weeks had a mean increase compared to baseline in fasting blood glucose of 2.3 mg/dL. Placebo-treated subjects (N=19) had a mean increase in fasting blood glucose compared to baseline of 0.34 mg/dL.

In an analysis of 7 controlled clinical studies, 2 of which were placebo-controlled, with treatment duration up to 12 weeks, olanzapine and fluoxetine was associated with a greater mean change in random glucose when compared to placebo (+8.65 mg/dL vs -3.86 mg/dL). The difference in mean changes between olanzapine and fluoxetine and placebo was greater in patients with evidence of glucose dysregulation at baseline (including those patients diagnosed with diabetes mellitus or related adverse reactions, patients treated with anti-diabetic agents, patients with a baseline random glucose level ≥200 mg/dL, or a baseline fasting glucose level ≥126 mg/dL). Olanzapine and fluoxetine-treated patients had a greater mean HbA_{1c} increase (baseline of 5.76% (median exposure 63 days), compared to a mean HbA_{1c} decrease of 0.04% in fluoxetine-treated subjects (median exposure 57 days) and a mean HbA_{1c} increase of 0.12% in olanzapine-treated patients (median exposure 56 days).

In an analysis of 6 controlled clinical studies, a larger proportion of olanzapine and fluoxetine-treated subjects had glycosuria (4.4%) compared to placebo-treated subjects (1.4%).

The mean change in nonfasting glucose in patients treated with at least 48 weeks was +5.9 mg/dL (N=425).

Table 2 shows short-term and long-term changes in random glucose levels from adult olanzapine and fluoxetine studies.

Table 2: Changes in Random Glucose Levels from Adult Olanzapine and Fluoxetine Studies

| Laboratory Analyte | Category

Additional Findings Observed in Clinical Studies

Sexual Dysfunction — In the pool of controlled olanzapine and fluoxetine studies in patients with bipolar depression, there were higher rates of the treatment-emergent adverse reactions decreased libido, anorgasmia, erectile dysfunction and abnormal ejaculation reported for olanzapine/fluoxetine combination than in the placebo group. The rates of decreased libido led to discontinuation in the olanzapine and fluoxetine group. In the controlled studies that contained a fluoxetine arm, the rates of decreased libido and abnormal ejaculation in the olanzapine and fluoxetine group were higher than the rates in the fluoxetine group. None of the differences were statistically significant.

Sexual dysfunction, including priapism, has been reported with all SSRIs. While it is difficult to know the precise risk of sexual dysfunction associated with the use of SSRIs, physicians should routinely inquire about such possible side effects.

There are no adequate and well-controlled studies examining sexual dysfunction with olanzapine and fluoxetine or fluoxetine treatment. Symptoms of sexual dysfunction occasionally persist after discontinuation of fluoxetine treatment.

Difference Among Dose Levels Observed in Other Olanzapine Clinical Trials

In a single 9-week randomized, double-blind, fixed-dose study comparing 10 (N=199), 20 (N=200), and 40 (N=200) mg/day of olanzapine with placebo (N=205), the incidence of patients who discontinued due to adverse reactions was comparable among 3 dose groups were observed for the following safety outcomes: weight gain, prolactin elevation, fatigue, and dizziness. Olanzapine treatment (mean dose 18 mg/day) was associated with a 4.4 kg (7.1%) mean weight gain, 3 kg) was observed with significant differences between 10 vs 40 mg/day. Incidence of treatment-emergent prolactin elevation >2 ng/mL (female) or >18.77 ng/mL (male) at any time during the trial (10 mg/day: 7.2%, 20 mg/day: 4.7%, 40 mg/day: 6.6%) did not differ significantly between 10 vs 40 mg/day and 20 vs 40 mg/day. Prolactin increase (>10 mg/day: 1.5%, 20 mg/day: 2.1%, 40 mg/day: 6.6%) with significant differences between 10 vs 40 and 20 vs 40 mg/day, and dizziness (10 mg/day: 2.6%, 20 mg/day: 1.6%, 40 mg/day: 6.6%) with significant differences between 20 vs 40 mg, was observed.

Other Adverse Reactions Observed in Clinical Studies

Following is a list of treatment-emergent adverse reactions reported by patients treated with olanzapine and fluoxetine in clinical trials. This listing is not intended to include reactions (1) already listed in previous tables or elsewhere in labeling, (2) for which adverse drug experience is not reasonably likely to be associated with the use of olanzapine and fluoxetine not considered to have significant clinical implications, or (5) which occurred at a rate equal to or less than placebo.

Reactions are classified by body system using the following definitions: frequent adverse reactions are those occurring in at least 1/100 patients; infrequent adverse reactions are those occurring in 1/100 to 1/1000 patients; and rare reactions are those occurring in fewer than 1/1000 patients.

Body as a Whole — *Frequent:* asthenia; *infrequent:* abdomen enlarged, allergic reaction; *Rare:* death[1]

Cardiovascular System — *Frequent:* vasodilation

Digestive System — *Frequent:* diarrhea; *infrequent:* gastritis, gastroenteritis, nausea and vomiting, peptic ulcer; *Rare:* gastrointestinal hemorrhage, intestinal obstruction, intestinal perforation, liver fatty deposit, pancreatitis

Hemic and Lymphatic System — *Frequent:* ecchymosis; *Infrequent:* anemia, thrombocytopenia; *Rare:* leukopenia, purpura

Metabolic and Nutritional — *Frequent:* generalized edema, weight loss; *Rare:* bilirubinemia, creatinine increased, gout

Musculoskeletal System — *Rare:* osteoporosis.

Nervous System — *Frequent:* amnesia; *Infrequent:* ataxia, buccoglossal syndrome, coma, depersonalization, dysarthria, emotional lability, hallucinations, hypertonia, hypesthesia, neuropathy, nystagmus; *Rare:* hyperkinesia, libido increased, withdrawal syndrome.

Respiratory System — *Infrequent:* epistaxis, yawn; *Rare:* laryngismus.

Skin and Appendages — *Infrequent:* alopecia, dry skin, pruritis; *Rare:* exfoliative dermatitis.

Special Senses — *Frequent:* taste perversion; *Infrequent:* abnormality of accommodation, dry eyes.

Urogenital System — *Frequent:* breast pain, menorrhagia, urinary frequency, urinary incontinence; *Infrequent:* amenorrhea[2], female lactation[2], hypomenorrhea[2], metrorrhagia[2], urinary retention, urinary urgency, urination impaired; *Rare:* breast engorgement[2].

† This term represents adverse event but does not meet the definition for adverse drug reactions. It is included here because of its seriousness.

‡ Adjusted for gender

Other Adverse Reactions Observed with Olanzapine or Fluoxetine Monotherapy

The following adverse reactions were not observed in olanzapine and fluoxetine-treated patients during premarketing clinical studies, but have been reported in olanzapine or fluoxetine monotherapy: neuroleptic malignant syndrome, seizures, cholestatic jaundice, diabetic coma, dysuria, eosinophilic pneumonia[1], erythema multiforme, esophageal ulcer, gynecological bleeding, hepatic failure/necrosis, hepatitis, hyponatremia, inappropriate antidiuretic hormone secretion, sudden unexpected death[1], sweating, and violent behaviors[1]. Random triglyceride levels of ≥1000 mg/dL have been reported.

† These terms represent serious adverse events but do not meet the definition for adverse drug reactions. They are included here because of their seriousness.

Children and Adolescent Patients (aged 10 to 17 years) with a Diagnosis of Bipolar Depression

The information below is derived from a single, 8-week, randomized, placebo-controlled clinical trial investigating olanzapine and fluoxetine for the treatment of bipolar depression in patients 10 to 17 years of age.

Adverse Reactions Associated with Discontinuation of Treatment in the single pediatric study — Overall, 14.1% of the 170 patients in the olanzapine and fluoxetine group discontinued treatment due to adverse reactions compared with 5.9% of the 85 patients for placebo. Adverse reactions leading to discontinuation associated with the use of olanzapine and fluoxetine (incidence of at least 1% for olanzapine and fluoxetine and greater than that for placebo) using MedDRA Preferred Term coding were weight increased (2.9%), suicidal ideation (1.8%), bipolar disorder (1.2%), and somnolence (1.2%) versus placebo patients which had 0% incidence of weight increased, bipolar disorder, and somnolence, and a 1.2% incidence of suicidal ideation.

Adverse Reactions Occurring at an Incidence of 2% or more and greater than placebo — **Table 17** enumerates the treatment-emergent adverse reactions associated with the use of olanzapine and fluoxetine (incidence of at least 2% for olanzapine and fluoxetine and twice or more than for placebo).

Table 17: Treatment-Emergent Adverse Reactions: Incidence in a 8-week randomized, double blind, placebo-controlled clinical trial in pediatric bipolar I depression

System Organ Class	Adverse Reaction	Percentage of Patients Reporting Event	
		Olanzapine and Fluoxetine (N=170)	Placebo (N=85)
Nervous system disorders	Somnolence[a]	24	2
	Tremor	9	1
Investigations	Weight Increased	20	1
	Blood triglycerides increased	7	0
	Blood cholesterol increased	4	0
	Hepatic Enzyme increased[b]	6	0
Gastrointestinal disorders	Dyspepsia	3	1
Metabolism and nutrition disorders	Increased appetite	17	1
Psychiatric disorders	Anxiety	3	1
	Restlessness	3	1
	Suicidal Ideation	2	1
Musculoskeletal and connective tissue disorders	Back pain	2	1
Injury, poisoning and procedural complications	Accidental overdose	3	1
Reproductive system and breast disorders	Dysmenorrhea	2	0

[a] Includes somnolence, sedation, and hypersomnia. No lethargy was reported.

[b] Includes alanine aminotransferase increased, aspartate aminotransferase increased, hepatic enzyme increased, liver function test abnormal, gamma-glutamyltransferase increased, and transaminases increased.

6.2 Vital Signs and Laboratory Studies

Adults:

Vital Signs — Tachycardia, bradycardia, and orthostatic hypotension have occurred in olanzapine and fluoxetine HCl-treated patients *[see Warnings and Precautions (5.11)]*. The mean standing pulse rate of olanzapine and fluoxetine-treated patients was reduced by 0-7 beats/min.

Laboratory Changes — In olanzapine and fluoxetine clinical studies (including treatment resistant depression, depressive episodes associated with Bipolar I Disorder, Major Depressive Disorder with psychosis, or sexual dysfunction) olanzapine and fluoxetine was not associated with statistically significantly greater frequencies for the following treatment-emergent findings in laboratory analytes (normal at baseline to abnormal at any time during the trial) compared to placebo: elevated creatinine (28% vs 5%), elevated urea nitrogen (3% vs 0.6%), elevated uric acid (3% vs 0.5%), low albumin (3% vs 0.3%), low bicarbonate (14% vs 9%), low hemoglobin (3% vs 0%), low inorganic phosphorus (1% vs 0%), low potassium (4% vs 2%), and low sodium (15% vs 4%).

As with olanzapine, asymptomatic elevations of hepatic aminotransferases [ALT, AST, and GGT] and alkaline phosphatase have been observed with olanzapine and fluoxetine. In the olanzapine and fluoxetine-controlled database, clinically significant ALT elevations (≥ 3 times the upper limit of normal [ULN] at baseline and to ≥3 times ULN) were observed in 5% (38/696) of patients exposed to olanzapine and fluoxetine compared with 0.5% (2/378) of placebo-treated patients. ALT elevations ≥5 times ULN were observed in 2% (13/701) of olanzapine-treated patients and 0.3% (1/378) of placebo-treated patients and 3% (11/701) of olanzapine-treated patients. No patient with elevated ALT values experienced jaundice or liver failure, or met the criteria for Hy's Rule. ALT values returned to normal, or were decreasing, at last follow-up in the majority of patients who either continued treatment with olanzapine and fluoxetine or discontinued olanzapine and fluoxetine.

Rare posttreatment reports of hepatitis have been received in patients treated with olanzapine. Very rare cases of cholestatic or mixed liver injury have also been reported in the postmarketing period in patients treated with olanzapine.

Caution should be exercised in patients with signs and symptoms of hepatic impairment, in patients with pre-existing conditions associated with limited hepatic functional reserve, and in patients who are being treated with potentially hepatotoxic drugs.

An increase in creatine phosphokinase has been reported very rarely in olanzapine and fluoxetine-treated patients and infrequently in clinical trials of olanzapine-treated patients.

QT Interval Prolongation — In patients treated with olanzapine and fluoxetine QTcF≥460 msec for males and QTcF≥470 msec for females has been reported in 2% of patients. The incidence of QTcF>500 msec reported with olanzapine and fluoxetine treatment in clinical trials has been rare and was not significantly different from the incidence associated with placebo. The mean increase in QTcF interval for olanzapine and fluoxetine-treated patients (5.17 msec) in the one clinical study directly comparing olanzapine and fluoxetine to placebo in adult patients was significantly greater than that for placebo-treated patients (-1.66 msec).

Children and Adolescent (aged 10 to 17 years):

In a single 8-week randomized, placebo-controlled clinical trial investigating olanzapine and fluoxetine for treatment of bipolar I depression in patients 10 to 17 years of age, the following was observed:

Vital Signs — In the olanzapine and fluoxetine treated patients compared with placebo-treated patients, the mean orthostatic blood pressure and standing pulse rate were not significantly different between treatment groups.

Body Weight: An increase in weight greater than or equal to 7% occurred in 52.4% of the olanzapine and fluoxetine group and 3.6% of the placebo group. Weight gain greater than or equal to 15% occurred in 14.1% of the olanzapine and fluoxetine group and none of the placebo group.

Laboratory Changes — Olanzapine and fluoxetine was associated with statistically significant greater frequencies for the following treatment-emergent findings in laboratory analytes (normal or low at baseline to abnormal at any time during the trial) compared to placebo: elevated ALT (45.9% vs 2.5%), elevated AST (33.7% vs 7.6%), high fasting total cholesterol (29.9% vs 8.2%); high fasting LDL cholesterol (19.7% vs 6.5%); high fasting triglycerides (52.3% vs 27.3%), and elevated prolactin (85% vs 36%). No patient with elevated hepatic enzyme values experienced jaundice or liver failure, or met the criteria for Hy's Rule. Five patients experienced an adverse event potentially associated with elevated prolactin: these events included galactorrhoea, galactostasis, and ovulation disorder.

QT Interval Prolongation — Olanzapine and Fluoxetine was associated with a statistically significantly greater mean increase in QT,F interval (8.2 msec [95% CI:6.2, 10.2]) compared with placebo. No patients developed QTc increases ≥60 msec or QTc ≥480 msec *[see Warnings and Precautions (5.20)]*.

6.3 Postmarketing Experience

The following adverse reactions have been identified during post-approval use of olanzapine and fluoxetine. Because these reactions are reported voluntarily from a population of uncertain size, it is difficult to reliably estimate their frequency or evaluate a causal relationship to drug exposure.

Adverse reactions reported since market introduction that were temporally (but not necessarily causally) related to olanzapine and fluoxetine therapy include the following: rhabdomyolysis and venous thromboembolic events (including pulmonary embolism and deep venous thrombosis).

7 DRUG INTERACTIONS

The risks of using olanzapine and fluoxetine in combination with other drugs have not been extensively evaluated in systematic studies. The drug-drug interactions sections of fluoxetine and olanzapine are applicable to olanzapine and fluoxetine. As with all drugs, the potential for interaction by a variety of mechanisms (e.g., pharmacodynamic, pharmacokinetic drug inhibition or enhancement, etc.) is a possibility. In evaluating individual cases, consideration should be given to using lower initial doses of the concomitantly administered drugs, using conservative titration schedules, and monitoring of clinical status *[see Clinical Pharmacology (12.3)]*.

7.1 Monoamine Oxidase Inhibitors (MAOIs)

[see Dosage and Administration (2.4, 2.5). Contraindications (4.1), and Warnings and Precautions (5.6)].

7.2 CNS Acting Drugs

Caution is advised if the concomitant administration of olanzapine and fluoxetine and other CNS-active drugs is required. In evaluating individual cases, consideration should be given to using lower initial doses of the concomitantly administered drugs, using conservative titration schedules, and monitoring of clinical status *[see Clinical Pharmacology (12.3)]*.

7.3 Serotonergic Drugs

[see Dosage and Administration (2.4, 2.5). Contraindications (4.1), and Warnings and Precautions (5.6)].

7.4 Drugs that Interfere with Hemostasis (e.g., NSAIDs, Aspirin, Warfarin)

Serotonin release by platelets plays an important role in hemostasis. Epidemiological studies of the case-control and cohort design that have demonstrated an association between use of psychotropic drugs that interfere with serotonin reuptake and the occurrence of upper gastrointestinal bleeding have also shown that concurrent use of an NSAID or aspirin may potentiate the risk of bleeding. Altered anticoagulant effects, including increased bleeding, have been reported when SNRIs or SSRIs are coadministered with warfarin *[see Warnings and Precautions (5.16)]*. Warfarin (2.0 mg given daily) did not affect olanzapine pharmacokinetics. Single doses of olanzapine did not affect the pharmacokinetics of warfarin. Patients receiving warfarin therapy should be carefully monitored when olanzapine and fluoxetine is started or discontinued.

7.5 Electroconvulsive Therapy (ECT)

There are no clinical studies establishing the benefit of the combined use of ECT and fluoxetine. There have been rare reports of prolonged seizures in patients on fluoxetine receiving ECT treatment *[see Warnings and Precautions (5.16)]*.

7.6 Potential for Other Drugs to Affect Olanzapine and Fluoxetine

Benzodiazepines — Coadministration of diazepam with olanzapine potentiated the orthostatic hypotension observed with olanzapine *[see Drug Interactions (7.7)]*.

Inducers of 1A2 — Carbamazepine therapy (200 mg BID) causes an approximate 50% increase in the clearance of olanzapine. This increase is likely due to the fact that carbamazepine is a potent inducer of CYP1A2 activity. Higher daily doses of carbamazepine may cause an even greater increase in olanzapine clearance *[see Drug Interactions (7.7)]*.

Alcohol — Ethanol (45 mg/70 kg single dose) did not have an effect on olanzapine pharmacokinetics *[see Drug Interactions (7.7)]*.

Inhibitors of CYP1A2 — Fluvoxamine decreases the clearance of olanzapine. This results in a mean increase in olanzapine C_{max} following fluvoxamine administration of 54% in female nonsmokers and 77% in male smokers. The mean increase in olanzapine AUC is 52% and 108%, respectively. Lower doses of the olanzapine component of olanzapine and fluoxetine should be considered in patients receiving concomitant treatment with fluvoxamine.

The Effect of Other Drugs on Olanzapine — Fluoxetine, an inhibitor of CYP2D6, decreases olanzapine clearance a small amount *[see Clinical Pharmacology (12.3)]*. Agents that induce CYP1A2 or glucuronyl transferase enzymes, such as omeprazole and rifampin, may cause an increase in olanzapine clearance. The effect of CYP1A2 inhibitors, such as fluvoxamine and some fluoroquinolone antibiotics, on olanzapine and fluoxetine has not been established. Although olanzapine is metabolized by multiple enzyme systems, induction or inhibition of a single enzyme may appreciably alter olanzapine clearance. Therefore, a dosage increase (for induction) or a dosage decrease (for inhibition) may need to be considered with specific drugs.

7.7 Potential for Olanzapine and Fluoxetine to Affect Other Drugs

Pimozide — Concomitant use of olanzapine and fluoxetine and pimozide is contraindicated. Pimozide can prolong the QT interval. Olanzapine and fluoxetine may prolong the QT interval. Coadministration of CYP2D6, CYP3A4, and CYP1A2 inhibitors and Fluoxetine can also prolong the QT interval. Clinical studies of pimozide with other antidepressants demonstrate an increase in drug interaction or QT,i prolongation. While a specific study with pimozide and olanzapine and fluoxetine has not been conducted, the potential for drug interactions or QT,i prolongation warrants restricting the concurrent use of pimozide and olanzapine and fluoxetine. *[see Contraindications (4.2), Warnings and Precautions (5.20) and Drug Interactions (7.8)]*.

Carbamazepine — Patients on stable doses of carbamazepine have developed elevated plasma anticonvulsant concentrations and clinical anticonvulsant toxicity following initiation of concomitant fluoxetine treatment.

Alcohol — The coadministration of ethanol with olanzapine and fluoxetine may potentiate sedation and orthostatic hypotension *[see Drug Interactions (7.6)]*.

Thioridazine — Thioridazine should not be administered with olanzapine and fluoxetine or administered within a minimum of 5 weeks after discontinuation of olanzapine and fluoxetine capsules, because of the risk of QT prolongation *[see Contraindications (4.2), Warnings and Precautions (5.20), and Drug Interactions (7.8)]*.

In a study of 19 healthy male subjects, which included 6 slow and 13 rapid hydroxylators of debrisoquin, a single 25 mg oral dose of thioridazine produced a 2.4-fold higher C_{max} and a 4.5-fold higher AUC for thioridazine and a 1.4-fold higher C_{max} and 2.0-fold higher AUC for its active metabolite. The rate of debrisoquin hydroxylation is known to depend on the level of CYP2D6 isozyme activity. Thus, this study suggests that drugs that inhibit CYP2D6, such as certain SSRIs, including fluoxetine, will increase the plasma levels of thioridazine *[see Contraindications (4.2)]*.

Thioridazine administration produces a dose-related prolongation of the QT_c interval, which is associated with serious ventricular arrhythmias, such as torsades de pointes-type arrhythmias and sudden death. This risk is expected to increase with fluoxetine administration. Coadministration of thioridazine and fluoxetine is contraindicated.

Due to the risk of serious ventricular arrhythmias and sudden death potentially associated with elevated thioridazine plasma levels, thioridazine should not be administered with fluoxetine or within a minimum of 5 weeks after fluoxetine has been discontinued *[see Contraindications (4.2)]*.

Tricyclic Antidepressants (TCAs) — Single doses of olanzapine did not affect the pharmacokinetics of imipramine or its active metabolite desipramine.

In 2 fluoxetine studies, previously stable plasma levels of imipramine and desipramine have increased >2-to 10-fold when fluoxetine has been administered in combination. This influence may persist for 3 weeks or longer after fluoxetine is discontinued. Thus, the dose of TCA may need to be reduced and plasma TCA concentrations may need to be monitored temporarily when olanzapine and fluoxetine is coadministered or has been recently discontinued *[see Warnings and Precautions (5.6), Drug Interactions (7.8)]*.

Antihypertensive Agents — Because of the potential for olanzapine to induce hypotension, olanzapine and fluoxetine may enhance the effects of certain antihypertensive agents *[see Warnings and Precautions (5.11)]*.

Levodopa and Dopamine Agonists — The olanzapine component of fluoxetine and fluoxetine may antagonize the effects of levodopa and dopamine agonists.

Benzodiazepines — The clearance of diazepam did not change when coadministered with olanzapine.

When concurrently administered with fluoxetine, the half-life of diazepam may be prolonged in some patients *[see Clinical Pharmacology (12.3)]*. Coadministration of alprazolam and fluoxetine may result in increased alprazolam plasma concentrations and in further psychomotor performance decrement due to increased alprazolam levels.

Clozapine — Elevation of blood levels of clozapine has been observed in patients receiving concomitant fluoxetine.

Haloperidol — Elevation of haloperidol blood levels and toxicity, including withdrawal symptoms following concomitant fluoxetine.

Drugs Metabolized by CYP2D6 — *In vitro* studies utilizing human liver microsomes suggest that olanzapine has little potential to inhibit CYP2D6. Thus, olanzapine is unlikely to cause clinically important drug interactions mediated by this enzyme.

Fluoxetine inhibits the activity of CYP2D6 and may make individuals with normal CYP2D6 metabolic activity resemble a poor metabolizer. Coadministration of fluoxetine and other drugs that are metabolized by CYP2D6, including certain antidepressants (e.g., TCAs), antipsychotics (e.g., phenothiazines and most atypicals), and antiarrhythmics (e.g., propafenone and flecainide), and others) should be approached with caution. The therapeutic activity of tricyclic antidepressants is predominantly metabolized by the CYP2D6 system and that have a relatively narrow therapeutic index should be initiated at the low end of the dose range if the patient is already receiving fluoxetine concurrently or has taken it in the previous 5 weeks. Fluoxetine is added to the treatment regimen of a patient already receiving a drug metabolized by CYP2D6, the need for a decreased dose of the original medication should be considered. Drugs with a narrow therapeutic index represent the greatest concern (including thioridazine, TCAs, carbamazepine, and antiarrhythmics) *[see Warnings and Precautions (5.6)]*.

Drugs Metabolized by CYP3A — *In vitro* studies utilizing human liver microsomes suggest that olanzapine has little potential to inhibit CYP3A. Thus, olanzapine is unlikely to cause clinically important drug interactions mediated by these enzymes.

In an *in vivo* interaction study involving the coadministration of fluoxetine with single doses of terfenadine (a CYP3A substrate), no increase in plasma terfenadine (parent drug) concentrations occurred with concomitant fluoxetine. In addition, *in vitro* studies have shown ketoconazole, a potent inhibitor of CYP3A activity, to be at least 100 times more potent than fluoxetine or norfluoxetine as an inhibitor of the metabolism of several substrates for this enzyme, including astemizole, cisapride, and midazolam. These data indicate that fluoxetine's extent of inhibition of CYP3A activity is not likely to be of clinical significance.

Effect of Olanzapine on Drugs Metabolized by Other CYP Enzymes — *In vitro* studies utilizing human liver microsomes suggest that olanzapine has little potential to inhibit CYP1A2, CYP2C9, and CYP2C19. Thus, olanzapine is unlikely to cause clinically important drug interactions mediated by these enzymes.

Lithium — Multiple doses of olanzapine did not influence the pharmacokinetics of lithium.

There have been reports of both increased and decreased lithium levels when lithium was used concomitantly with fluoxetine. Cases of lithium toxicity and increased lithium levels have been reported in patients receiving lithium concomitantly with fluoxetine. Lithium levels should be monitored in patients taking olanzapine and fluoxetine concomitantly with lithium *[see Warnings and Precautions (5.5)]*.

Drugs Tightly Bound to Plasma Proteins — *The in vitro* binding of olanzapine and fluoxetine to human plasma proteins is similar to the individual components. The interaction between olanzapine and fluoxetine and other highly protein-bound drugs has not been fully evaluated. Because fluoxetine is highly protein bound, administration of fluoxetine to a patient taking another drug that is tightly bound to protein (e.g., Coumadin, digitoxin) may cause a shift in plasma concentrations potentially resulting in an adverse effect. Conversely, displacement of drug, have been reported when SNRIs or SSRIs are coadministered with warfarin *[see Warnings and Precautions (5.16)]*. Warfarin (2.0 mg given daily) did not affect olanzapine pharmacokinetics. Single doses of olanzapine did not affect the pharmacokinetics of warfarin. Patients receiving warfarin therapy should be carefully monitored when olanzapine and fluoxetine is started or discontinued.

7.5 Electroconvulsive Therapy (ECT)

There are no clinical studies establishing the benefit of the combined use of ECT and fluoxetine. There have been rare reports of prolonged seizures in patients on fluoxetine receiving ECT treatment *[see Warnings and Precautions (5.16)]*.

7.6 Potential for Other Drugs to Affect Olanzapine and Fluoxetine

Benzodiazepines — Coadministration of diazepam with olanzapine potentiated the orthostatic hypotension observed with olanzapine *[see Drug Interactions (7.7)]*.

Inducers of 1A2 — Carbamazepine therapy (200 mg BID) causes an approximate 50% increase in the clearance of olanzapine. This increase is likely due to the fact that carbamazepine is a potent inducer of CYP1A2 activity. Higher daily doses of carbamazepine may cause an even greater increase in olanzapine clearance *[see Drug Interactions (7.7)]*.

Alcohol — Ethanol (45 mg/70 kg single dose) did not have an effect on olanzapine pharmacokinetics *[see Drug Interactions (7.7)]*.

Inhibitors of CYP1A2 — Fluvoxamine decreases the clearance of olanzapine. This results in a mean increase in olanzapine C_{max} following fluvoxamine administration of 54% in female nonsmokers and 77% in male smokers. The mean increase in olanzapine AUC is 52% and 108%, respectively. Lower doses of the olanzapine component of olanzapine and fluoxetine should be considered in patients receiving concomitant treatment with fluvoxamine.

The Effect of Other Drugs on Olanzapine — Fluoxetine, an inhibitor of CYP2D6, decreases olanzapine clearance a small amount *[see Clinical Pharmacology (12.3)]*. Agents that induce CYP1A2 or glucuronyl transferase enzymes, such as omeprazole and rifampin, may cause an increase in olanzapine clearance. The effect of CYP1A2 inhibitors, such as fluvoxamine and some fluoroquinolone antibiotics, on olanzapine and fluoxetine has not been established. Although olanzapine is metabolized by multiple enzyme systems, induction or inhibition of a single enzyme may appreciably alter olanzapine clearance. Therefore, a dosage increase (for induction) or a dosage decrease (for inhibition) may need to be considered with specific drugs.

7.7 Potential for Olanzapine and Fluoxetine to Affect Other Drugs

Pimozide — Concomitant use of olanzapine and fluoxetine and pimozide is contraindicated. Pimozide can prolong the QT interval. Olanzapine and fluoxetine may prolong the QT interval. Coadministration of CYP2D6, CYP3A4, and CYP1A2 inhibitors and Fluoxetine can also prolong the QT interval. Clinical studies of pimozide with other antidepressants demonstrate an increase in drug interaction or QT,i prolongation. While a specific study with pimozide and olanzapine and fluoxetine has not been conducted, the potential for drug interactions or QT,i prolongation warrants restricting the concurrent use of pimozide and olanzapine and fluoxetine. *[see Contraindications (4.2), Warnings and Precautions (5.20) and Drug Interactions (7.8)]*.

Carbamazepine — Patients on stable doses of carbamazepine have developed elevated plasma anticonvulsant concentrations and clinical anticonvulsant toxicity following initiation of concomitant fluoxetine treatment.

Alcohol — The coadministration of ethanol with olanzapine and fluoxetine may potentiate sedation and orthostatic hypotension *[see Drug Interactions (7.6)]*.

Thioridazine — Thioridazine should not be administered with olanzapine and fluoxetine or administered within a minimum of 5 weeks after discontinuation of olanzapine and fluoxetine capsules, because of the risk of QT prolongation *[see Contraindications (4.2), Warnings and Precautions (5.20), and Drug Interactions (7.8)]*.

In a study of 19 healthy male subjects, which included 6 slow and 13 rapid hydroxylators of debrisoquin, a single 25 mg oral dose of thioridazine produced a 2.4-fold higher C_{max} and a 4.5-fold higher AUC for thioridazine and a 1.4-fold higher C_{max} and 2.0-fold higher AUC for its active metabolite. The rate of debrisoquin hydroxylation is known to depend on the level of CYP2D6 isozyme activity. Thus, this study suggests that drugs that inhibit CYP2D6, such as certain SSRIs, including fluoxetine, will increase the plasma levels of thioridazine *[see Contraindications (4.2)]*.

Thioridazine administration produces a dose-related prolongation of the QT_c interval, which is associated with serious ventricular arrhythmias, such as torsades de pointes-type arrhythmias and sudden death. This risk is expected to increase with fluoxetine administration. Coadministration of thioridazine and fluoxetine is contraindicated.

Due to the risk of serious ventricular arrhythmias and sudden death potentially associated with elevated thioridazine plasma levels, thioridazine should not be administered with fluoxetine or within a minimum of 5 weeks after fluoxetine has been discontinued *[see Contraindications (4.2)]*.

Tricyclic Antidepressants (TCAs) — Single doses of olanzapine did not affect the pharmacokinetics of imipramine or its active metabolite desipramine.

In 2 fluoxetine studies, previously stable plasma levels of imipramine and desipramine have increased >2-to 10-fold when fluoxetine has been administered in combination. This influence may persist for 3 weeks or longer after fluoxetine is discontinued. Thus, the dose of TCA may need to be reduced and plasma TCA concentrations may need to be monitored temporarily when olanzapine and fluoxetine is coadministered or has been recently discontinued *[see Warnings and Precautions (5.6), Drug Interactions (7.8)]*.

Antihypertensive Agents — Because of the potential for olanzapine to induce hypotension, olanzapine and fluoxetine may enhance the effects of certain antihypertensive agents *[see Warnings and Precautions (5.11)]*.

Levodopa and Dopamine Agonists — The olanzapine component of fluoxetine and fluoxetine may antagonize the effects of levodopa and dopamine agonists.

Benzodiazepines — The clearance of diazepam did not change when coadministered with olanzapine.

When concurrently administered with fluoxetine, the half-life of diazepam may be prolonged in some patients *[see Clinical Pharmacology (12.3)]*. Coadministration of alprazolam and fluoxetine may result in increased alprazolam plasma concentrations and in further psychomotor performance decrement due to increased alprazolam levels.

Clozapine — Elevation of blood levels of clozapine has been observed in patients receiving concomitant fluoxetine.

Haloperidol — Elevation of haloperidol blood levels and toxicity, including withdrawal symptoms following concomitant fluoxetine.

Drugs Metabolized by CYP2D6 — *In vitro* studies utilizing human liver microsomes suggest that olanzapine has little potential to inhibit CYP2D6. Thus, olanzapine is unlikely to cause clinically important drug interactions mediated by this enzyme.

Fluoxetine inhibits the activity of CYP2D6 and may make individuals with normal CYP2D6 metabolic activity resemble a poor metabolizer. Coadministration of fluoxetine and other drugs that are metabolized by CYP2D6, including certain antidepressants (e.g., TCAs), antipsychotics (e.g., phenothiazines and most atypicals), and antiarrhythmics (e.g., propafenone and flecainide), and others) should be approached with caution. The therapeutic activity of tricyclic antidepressants is predominantly metabolized by the CYP2D6 system and that have a relatively narrow therapeutic index should be initiated at the low end of the dose range if the patient is already receiving fluoxetine concurrently or has taken it in the previous 5 weeks. If fluoxetine is added to the treatment regimen of a patient already receiving a drug metabolized by CYP2D6, the need for a decreased dose of the original medication should be considered. Drugs with a narrow therapeutic index represent the greatest concern (including thioridazine, TCAs, carbamazepine, and antiarrhythmics) *[see Warnings and Precautions (5.6)]*.

Drugs Metabolized by CYP3A — *In vitro* studies utilizing human liver microsomes suggest that olanzapine has little potential to inhibit CYP3A. Thus, olanzapine is unlikely to cause clinically important drug interactions mediated by these enzymes.

In an *in vivo* interaction study involving the coadministration of fluoxetine with single doses of terfenadine (a CYP3A substrate), no increase in plasma terfenadine (parent drug) concentrations occurred with concomitant fluoxetine. In addition, *in vitro* studies have shown ketoconazole, a potent inhibitor of CYP3A activity, to be at least 100 times more potent than fluoxetine or norfluoxetine as an inhibitor of the metabolism of several substrates for this enzyme, including astemizole, cisapride, and midazolam. These data indicate that fluoxetine's extent of inhibition of CYP3A activity is not likely to be of clinical significance.

Effect of Olanzapine on Drugs Metabolized by Other CYP Enzymes — *In vitro* studies utilizing human liver microsomes suggest that olanzapine has little potential to inhibit CYP1A2, CYP2C9, and CYP2C19. Thus, olanzapine is unlikely to cause clinically important drug interactions mediated by these enzymes.

Lithium — Multiple doses of olanzapine did not influence the pharmacokinetics of lithium.

There have been reports of both increased and decreased lithium levels when lithium was used concomitantly with fluoxetine. Cases of lithium toxicity and increased lithium levels have been reported in patients receiving lithium concomitantly with fluoxetine. Lithium levels should be monitored in patients taking olanzapine and fluoxetine concomitantly with lithium *[see Warnings and Precautions (5.5)]*.

Drugs Tightly Bound to Plasma Proteins — *The in vitro* binding of olanzapine and fluoxetine to human plasma proteins is similar to the individual components. The interaction between olanzapine and fluoxetine and other highly protein-bound drugs has not been fully evaluated. Because fluoxetine is highly protein bound, administration of fluoxetine to a patient taking another drug that is tightly bound to protein (e.g., Coumadin, digitoxin) may cause a shift in plasma concentrations potentially resulting in an adverse effect. Conversely, displacement of drug, may have been reported with SNRIs or SSRIs are coadministered with warfarin *[see Warnings and Precautions (5.16)]*. Warfarin (2.0 mg given daily) did not affect olanzapine pharmacokinetics. Single doses of olanzapine did not affect the pharmacokinetics of warfarin. Patients receiving warfarin therapy should be carefully monitored when olanzapine and fluoxetine is started or discontinued.

7.5 Electroconvulsive Therapy (ECT)

There are no clinical studies establishing the benefit of the combined use of ECT and fluoxetine. There have been rare reports of prolonged seizures in patients on fluoxetine receiving ECT treatment *[see Warnings and Precautions (5.16)]*.

7.6 Potential for Other Drugs to Affect Olanzapine and Fluoxetine

Benzodiazepines — Coadministration of diazepam with olanzapine potentiated the orthostatic hypotension observed with olanzapine *[see Drug Interactions (7.7)]*.

Inducers of 1A2 — Carbamazepine therapy (200 mg BID) causes an approximate 50% increase in the clearance of olanzapine. This increase is likely due to the fact that carbamazepine is a potent inducer of CYP1A2 activity. Higher daily doses of carbamazepine may cause an even greater increase in olanzapine clearance *[see Drug Interactions (7.7)]*.

Alcohol — Ethanol (45 mg/70 kg single dose) did not have an effect on olanzapine pharmacokinetics *[see Drug Interactions (7.7)]*.

Inhibitors of CYP1A2 — Fluvoxamine decreases the clearance of olanzapine. This results in a mean increase in olanzapine C_{max} following fluvoxamine administration of 54% in female nonsmokers and 77% in male smokers. The mean increase in olanzapine AUC is 52% and 108%, respectively. Lower doses of the olanzapine component of olanzapine and fluoxetine should be considered in patients receiving concomitant treatment with fluvoxamine.

The Effect of Other Drugs on Olanzapine — Fluoxetine, an inhibitor of CYP2D6, decreases olanzapine clearance a small amount *[see Clinical Pharmacology (12.3)]*. Agents that induce CYP1A2 or glucuronyl transferase enzymes, such as omeprazole and rifampin, may cause an increase in olanzapine clearance. The effect of CYP1A2 inhibitors, such as fluvoxamine and some fluoroquinolone antibiotics, on olanzapine and fluoxetine has not been established. Although olanzapine is metabolized by multiple enzyme systems, induction or inhibition of a single enzyme may appreciably alter olanzapine clearance. Therefore, a dosage increase (for induction) or a dosage decrease (for inhibition) may need to be considered with specific drugs.

7.7 Potential for Olanzapine and Fluoxetine to Affect Other Drugs

Pimozide — Concomitant use of olanzapine and fluoxetine and pimozide is contraindicated. Pimozide can prolong the QT interval. Olanzapine and fluoxetine may prolong the QT interval. Coadministration of CYP2D6, CYP3A4, and CYP1A2 inhibitors and Fluoxetine can also prolong the QT interval. Clinical studies of pimozide with other antidepressants demonstrate an increase in drug interaction or QT,i prolongation. While a specific study with pimozide and olanzapine and fluoxetine has not been conducted, the potential for drug interactions or QT,i prolongation warrants restricting the concurrent use of pimozide and olanzapine and fluoxetine. *[see Contraindications (4.2), Warnings and Precautions (5.20) and Drug Interactions (7.8)]*.

Carbamazepine — Patients on stable doses of carbamazepine have developed elevated plasma anticonvulsant concentrations and clinical anticonvulsant toxicity following initiation of concomitant fluoxetine treatment.

Alcohol — The coadministration of ethanol with olanzapine and fluoxetine may potentiate sedation and orthostatic hypotension *[see Drug Interactions (7.6)]*.

Thioridazine — Thioridazine should not be administered with ol